

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use INVOKAMET®/INVOKAMET XR safely and effectively.

See full prescribing information for INVOKAMET/INVOKAMET XR.

INVOKAMET (canagliflozin and metformin hydrochloride) tablets, for oral use
INVOKAMET XR (canagliflozin and metformin hydrochloride) extended-release tablets, for oral use

Initial U.S. Approval: 2014

WARNING: LACTIC ACIDOSIS and LOWER LIMB AMPUTATION *See full prescribing information for complete boxed warning.*

Lactic Acidosis

- Postmarketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. Symptoms included malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Laboratory abnormalities included elevated blood lactate levels, anion gap acidosis, increased lactate/pyruvate ratio; and metformin plasma levels generally >5 mcg/mL. (5.1)
- Risk factors include renal impairment, concomitant use of certain drugs, age >65 years old, radiological studies with contrast, surgery and other procedures, hypoxic states, excessive alcohol intake, and hepatic impairment. Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high risk groups are provided in the Full Prescribing Information. (5.1)
- If lactic acidosis is suspected, discontinue INVOKAMET/INVOKAMET XR and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended. (5.1)

Risk of Lower Limb Amputation

- In patients with type 2 diabetes who have established cardiovascular disease (CVD) or at risk for CVD, canagliflozin, a component of INVOKAMET/INVOKAMET XR, has been associated with lower limb amputations, most frequently of the toe and midfoot; some also involved the leg. (5.2)
- Before initiating, consider factors that may increase the risk of amputation. Monitor patients receiving INVOKAMET/INVOKAMET XR for infections or ulcers of the lower limbs, and discontinue if these occur. (5.2)

-----RECENT MAJOR CHANGES-----

Indications and Usage (1)	10/2018
Warnings and Precautions (5.2, 5.7, 5.8, 5.12)	10/2018
Warnings and Precautions (5.6, 5.14)	Removal 10/2018

-----INDICATIONS AND USAGE-----

INVOKAMET and INVOKAMET XR are a combination of canagliflozin, a sodium-glucose co-transporter 2 (SGLT2) inhibitor, and metformin hydrochloride (HCl), a biguanide, indicated:

- as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus when treatment with both canagliflozin and metformin HCl is appropriate (1)
- Canagliflozin is indicated to reduce the risk of major adverse cardiovascular events in adults with type 2 diabetes mellitus and established cardiovascular disease (1)

Limitations of Use:

- Not for treatment of type 1 diabetes mellitus or diabetic ketoacidosis (1)

-----DOSAGE AND ADMINISTRATION-----

- Individualize starting dose based on the patient's current regimen (2.2, 2.3)
- INVOKAMET: one tablet, twice daily with meals, recommended starting dose of canagliflozin is 50 mg twice daily and metformin HCl 500 mg twice daily (2.2)
- INVOKAMET XR: two tablets, once daily with the morning meal. Swallow whole. Never crush, cut, or chew (2.2)
- Canagliflozin dose can be increased to a total daily dose of 300 mg in patients tolerating 100 mg who have an eGFR of 60 mL/min/1.73 m² or greater and require additional glycemic control. Do not exceed a total daily canagliflozin dose of 300 mg (2.2)
- Gradually escalate metformin HCl dose to reduce the gastrointestinal side effects while not exceeding a total daily dose of 2000 mg (2.1)
- Assess renal function before initiating and periodically thereafter (2.1, 2.3)
- Limit the dose of the canagliflozin component to two 50 mg tablets daily in patients with an eGFR of 45 to less than 60 mL/min/1.73 m² (2.4)
- May need to be discontinued at time of, or prior to, iodinated contrast imaging procedures (2.6)

-----DOSAGE FORMS AND STRENGTHS-----

INVOKAMET tablets:

- Canagliflozin 50 mg and metformin hydrochloride 500 mg
- Canagliflozin 50 mg and metformin hydrochloride 1,000 mg
- Canagliflozin 150 mg and metformin hydrochloride 500 mg
- Canagliflozin 150 mg and metformin hydrochloride 1,000 mg (3)

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

INVOKAMET XR extended-release tablets:

- Canagliflozin 50 mg and metformin hydrochloride 500 mg
- Canagliflozin 50 mg and metformin hydrochloride 1,000 mg
- Canagliflozin 150 mg and metformin hydrochloride 500 mg
- Canagliflozin 150 mg and metformin hydrochloride 1,000 mg (3)

-----CONTRAINDICATIONS-----

- Moderate to severe renal impairment (eGFR below 45 mL/min/1.73 m²), end stage renal disease or dialysis (4, 5.1, 5.5)
- Metabolic acidosis, including diabetic ketoacidosis (4, 5.1)
- Serious hypersensitivity reaction to canagliflozin or metformin (4, 5.10)

-----WARNINGS AND PRECAUTIONS-----

- **Hypotension:** Before initiating, assess volume status and correct hypovolemia in patients with renal impairment, the elderly, in patients with low systolic blood pressure, or on diuretics, ACEi, or ARB. Monitor for signs and symptoms during therapy (5.3)
- **Ketoacidosis:** Assess patients who present with signs and symptoms of metabolic acidosis for ketoacidosis, regardless of blood glucose level. If suspected, discontinue, evaluate and treat promptly. Before initiating, consider risk factors for ketoacidosis. Patients may require monitoring and temporary discontinuation in clinical situations known to predispose to ketoacidosis (5.4)
- **Acute kidney injury:** Consider temporarily discontinuing in settings of reduced oral intake or fluid losses. If acute kidney injury occurs, discontinue and promptly treat. Monitor renal function during therapy (5.5)
- **Ursepsis and pyelonephritis:** Evaluate patients for signs and symptoms of urinary tract infections and treat promptly, if indicated (5.6)
- **Hypoglycemia:** Consider a lower dose of insulin or insulin secretagogue to reduce the risk of hypoglycemia when used in combination (5.7)
- **Necrotizing fasciitis of the perineum (Fournier's gangrene):** Serious, life-threatening cases have occurred in both females and males. Assess patients presenting with pain or tenderness, erythema, or swelling in the genital or perineal area, along with fever or malaise. If suspected, institute prompt treatment (5.8)
- **Genital mycotic infections:** Monitor and treat if indicated (5.9)
- **Hypersensitivity reactions:** Discontinue and monitor until signs and symptoms resolve (5.10)
- **Bone fracture:** Consider factors that contribute to fracture risk before initiating INVOKAMET/INVOKAMET XR (5.11)
- **Vitamin B₁₂ deficiency:** Metformin may lower vitamin B₁₂ levels. Measure hematological parameters annually and vitamin B₁₂ at 2- to 3-year intervals and manage any abnormalities (5.12)
- **Increased LDL-C:** Monitor LDL-C and treat if appropriate (5.13)

-----ADVERSE REACTIONS-----

- Most common adverse reactions associated with canagliflozin (5% or greater incidence): female genital mycotic infections, urinary tract infection, and increased urination (6.1)
- Most common adverse reactions associated with metformin (5% or greater incidence) are diarrhea, nausea, vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Janssen Pharmaceuticals, Inc. at 1-800-526-7736 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

-----DRUG INTERACTIONS-----

- **Carbonic anhydrase inhibitors** may increase risk of lactic acidosis. Consider more frequent monitoring (7.1)
- **Drugs that reduce metformin clearance** (such as ranolazine, vandetanib, dolutegravir, and cimetidine) may increase the accumulation of metformin. Consider the benefits and risks of concomitant use (7.1)
- **Alcohol:** Warn patients against excessive intake (7.1)
- **UGT inducers** (e.g., rifampin): Canagliflozin exposure is reduced. Consider increasing canagliflozin from a total daily dose of 100 mg to a total daily dose of 300 mg (2.5, 7.2)
- **Digoxin:** Monitor digoxin levels (7.2)

-----USE IN SPECIFIC POPULATIONS-----

- **Pregnancy:** Advise females of the potential risk to a fetus especially during the second and third trimesters (8.1)
- **Lactation:** Not recommended when breastfeeding (8.2)
- **Females and Males of Reproductive Potential:** Advise premenopausal females of the potential for an unintended pregnancy (8.3).
- **Geriatrics:** Higher incidence of adverse reactions related to reduced intravascular volume. Assess renal function more frequently (5.3, 6.1, 8.5)
- **Renal impairment:** Higher incidence of adverse reactions related to reduced intravascular volume and renal function (2.4, 5.5, 8.6)
- **Hepatic Impairment:** Avoid use in patients with hepatic impairment (8.7)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide.

Revised: 10/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: LACTIC ACIDOSIS AND LOWER LIMB AMPUTATION

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Prior to Initiation of INVOKAMET/INVOKAMET XR
- 2.2 Dosage Overview
- 2.3 Starting Dose Based on Patient's Current Regimen
- 2.4 Recommended Dosage for Patients with Renal Impairment
- 2.5 Concomitant Use with UDP-Glucuronosyltransferase Enzyme Inducers
- 2.6 Discontinuation for Iodinated Contrast Imaging Procedures

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Lactic Acidosis
- 5.2 Lower Limb Amputation
- 5.3 Hypotension
- 5.4 Ketoacidosis
- 5.5 Acute Kidney Injury
- 5.6 Urosepsis and Pyelonephritis
- 5.7 Hypoglycemia with Concomitant Use of Sulfonylurea or Insulin
- 5.8 Necrotizing Fasciitis of the Perineum (Fournier's Gangrene)
- 5.9 Genital Mycotic Infections
- 5.10 Hypersensitivity Reactions
- 5.11 Bone Fracture
- 5.12 Vitamin B₁₂ Levels
- 5.13 Increases in Low-Density Lipoprotein (LDL-C)

6 ADVERSE REACTIONS

- 6.1 Clinical Studies Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

- 7.1 Drug Interactions with Metformin
- 7.2 Drug Interactions with Canagliflozin

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.3 Females and Males of Reproductive Potential
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Glycemic Control Trials in Adults with Type 2 Diabetes Mellitus
- 14.2 Canagliflozin Cardiovascular Outcomes in Patients with Type 2 Diabetes Mellitus and Atherosclerotic Cardiovascular Disease

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

*Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: LACTIC ACIDOSIS and LOWER LIMB AMPUTATION

Lactic Acidosis

- Post-marketing cases of metformin-associated lactic acidosis have resulted in death, hypothermia, hypotension, and resistant bradyarrhythmias. The onset of metformin-associated lactic acidosis is often subtle, accompanied only by nonspecific symptoms such as malaise, myalgias, respiratory distress, somnolence, and abdominal pain. Metformin-associated lactic acidosis was characterized by elevated blood lactate levels (> 5 mmol/Liter), anion gap acidosis (without evidence of ketonuria or ketonemia), an increased lactate/pyruvate ratio; and metformin plasma levels generally >5 mcg/mL [see *Warnings and Precautions (5.1)*].
- Risk factors for metformin-associated lactic acidosis include renal impairment, concomitant use of certain drugs (e.g., carbonic anhydrase inhibitors such as topiramate), age 65 years old or greater, having a radiological study with contrast, surgery and other procedures, hypoxic states (e.g., acute congestive heart failure), excessive alcohol intake, and hepatic impairment [see *Warnings and Precautions (5.1)*].
- Steps to reduce the risk of and manage metformin-associated lactic acidosis in these high risk groups are provided in the full prescribing information [see *Dosage and Administration (2.2, 2.3)*, *Contraindications (4)*, *Warnings and Precautions (5.1)*, *Drug Interactions (7)*, and *Use in Specific Populations (8.6, 8.7)*].
- If metformin-associated lactic acidosis is suspected, immediately discontinue INVOKAMET/INVOKAMET XR and institute general supportive measures in a hospital setting. Prompt hemodialysis is recommended [see *Warnings and Precautions (5.1)*].

Risk of Lower Limb Amputation

- An approximately 2-fold increased risk of lower limb amputations associated with canagliflozin, a component of INVOKAMET/INVOKAMET XR, was observed in CANVAS and CANVAS-R, two large, randomized, placebo-controlled trials in patients with type 2 diabetes who had established cardiovascular disease (CVD) or were at risk for CVD [see *Warnings and Precautions (5.2)*].
- Amputations of the toe and midfoot were most frequent; however, amputations involving the leg were also observed. Some patients had multiple amputations, some involving both limbs [see *Warnings and Precautions (5.2)*].
- Before initiating, consider factors that may increase the risk of amputation, such as a history of prior amputation, peripheral vascular disease, neuropathy, and diabetic foot ulcers [see *Warnings and Precautions (5.2)*].
- Monitor patients receiving INVOKAMET/INVOKAMET XR for infection, new pain or tenderness, sores or ulcers involving the lower limbs, and discontinue if these complications occur [see *Warnings and Precautions (5.2)*].

1 INDICATIONS AND USAGE

INVOKAMET and INVOKAMET XR are a combination of canagliflozin and metformin hydrochloride (HCl) indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes mellitus when treatment with both canagliflozin and metformin HCl is appropriate.

Canagliflozin is indicated to reduce the risk of major adverse cardiovascular events (cardiovascular death, nonfatal myocardial infarction and nonfatal stroke) in adults with type 2 diabetes mellitus and established cardiovascular disease (CVD) [see *Clinical Studies (14.2)*]. However, the effectiveness of INVOKAMET/INVOKAMET XR on reducing the risk of major cardiovascular events in adults with type 2 diabetes and cardiovascular disease has not been established.

Limitations of Use

Not recommended in patients with type 1 diabetes mellitus or for the treatment of diabetic ketoacidosis.

2 DOSAGE AND ADMINISTRATION

2.1 Prior to Initiation of INVOKAMET/INVOKAMET XR

- Assess renal function before initiating INVOKAMET/INVOKAMET XR and periodically thereafter [see *Warnings and Precautions (5.1, 5.5)*].
- In patients with volume depletion not previously treated with canagliflozin, normalize volume status before initiating INVOKAMET/INVOKAMET XR [see *Warnings and Precautions (5.3)*, *Use in Specific Populations (8.5, 8.6)*].

2.2 Dosage Overview

- INVOKAMET dosing is one tablet, twice daily with meals.
- INVOKAMET XR dosing is two tablets, once daily with the morning meal. Swallow each tablet whole and never crush, cut, or chew.
- The starting dose of INVOKAMET/INVOKAMET XR is based on the patient's current regimen [see *Dosage and Administration (2.2)*].
- The maximum recommended daily dose is canagliflozin 300 mg and metformin HCl 2,000 mg in patients with an estimated glomerular filtration rate (eGFR) of 60 mL/min/1.73 m² or greater. For patients with renal impairment [see *Dosage and Administration 2.4*].

2.3 Starting Dose Based on Patient's Current Regimen

Individualize the starting dose of INVOKAMET/INVOKAMET XR based on the patient's current regimen:

Patients Currently not Treated with Either Canagliflozin or Metformin HCl

- INVOKAMET: One tablet, twice daily, each tablet containing canagliflozin 50 mg and metformin HCl 500 mg;
- INVOKAMET XR: Two tablets, once daily, each tablet containing canagliflozin 50 mg and metformin HCl 500 mg.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Patients on Metformin

- INVOKAMET: One tablet, twice daily, where the total daily dose contains canagliflozin 100 mg and the same, or nearest appropriate, daily dose of metformin HCl;
- INVOKAMET XR: Two tablets, once daily, where the total daily dose contains canagliflozin 100 mg and the patient's same, or nearest appropriate, daily dose of metformin HCl.

Patients on Canagliflozin

- INVOKAMET: One tablet, twice daily, where the total daily dose contains the same daily dose of canagliflozin and metformin HCl 1,000 mg;
- INVOKAMET XR: Two tablets, once daily, where the total daily dose contains the patient's same daily dose of canagliflozin and metformin HCl 1,000 mg.

Patients Already Treated with Canagliflozin and Metformin

- INVOKAMET: One tablet, twice daily, where the total daily dose contains the same daily dose of canagliflozin and the same, or nearest appropriate, daily dose of metformin HCl;
- INVOKAMET XR: Two tablets, once daily, where the total daily dose contains the same daily dose of canagliflozin and the same, or nearest appropriate, daily dose of metformin HCl.

Patients with eGFR of 60 mL/min/1.73 m² or Greater and Require Additional Glycemic Control

- INVOKAMET: For patients tolerating canagliflozin 50 mg twice daily, increase the dose of canagliflozin to 150 mg twice daily, with gradual metformin HCl dose escalation based on tolerability [see *Adverse Reactions (6.1)*];
- INVOKAMET XR: For patients tolerating a daily dose of canagliflozin 100 mg, increase the dose to 300 mg once daily, with gradual metformin HCl dose escalation based on tolerability [see *Adverse Reactions (6.1)*].

Patients on Evening Dose of Metformin HCl Extended-Release

Patients taking an evening dose of metformin HCl extended-release should skip their last dose before starting INVOKAMET/INVOKAMET XR the following morning.

2.4 Recommended Dosage for Patients with Renal Impairment

- INVOKAMET/INVOKAMET XR is contraindicated in patients with an eGFR below 45 mL/min/1.73 m² [see *Contraindications (4) and Warnings and Precautions (5.1, 5.5)*].
- Limit the dose of the canagliflozin component of INVOKAMET/INVOKAMET XR to 100 mg (two 50 mg tablets) daily in patients with moderate renal impairment with an eGFR of 45 to less than 60 mL/min/1.73 m².

2.5 Concomitant Use with UDP-Glucuronosyltransferase Enzyme Inducers

If an inducer of UDP-Glucuronosyltransferase (UGT) (e.g., rifampin, phenytoin, phenobarbital, ritonavir) is co-administered with INVOKAMET/INVOKAMET XR, consider increasing the total daily dose of canagliflozin to 300 mg in patients currently tolerating a total daily dose of canagliflozin 100 mg who have an eGFR of 60 mL/min/1.73 m² or greater and require additional glycemic control [see *Drug Interactions (7.2)*].

Consider another antihyperglycemic agent in patients with an eGFR of 45 to less than 60 mL/min/1.73 m² receiving concurrent therapy with a UGT inducer.

2.6 Discontinuation for Iodinated Contrast Imaging Procedures

Discontinue INVOKAMET/INVOKAMET XR at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR of 45 to less than 60 mL/min/1.73 m²; in patients with a history of liver disease, alcoholism or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure; restart INVOKAMET/INVOKAMET XR if renal function is stable [see *Warnings and Precautions (5.1)*].

3 DOSAGE FORMS AND STRENGTHS

INVOKAMET (canagliflozin and metformin HCl) tablets are available as follows:

Canagliflozin Strength	Metformin HCl Strength	Color/Shape	Tablet Identifiers*
50 mg	500 mg	white/capsule-shaped	CM 155
50 mg	1,000 mg	beige/capsule-shaped	CM 551
150 mg	500 mg	yellow/capsule-shaped	CM 215
150 mg	1,000 mg	purple/capsule-shaped	CM 611

* Embossing appears on both sides of tablet.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

INVOKAMET XR (canagliflozin and metformin HCl) extended-release tablets are available as follows:

Canagliflozin Strength	Metformin HCl Strength	Color/Shape	Tablet Identifiers*
50 mg	500 mg	almost white to light orange/oblong, biconvex	CM1
50 mg	1,000 mg	pink/oblong, biconvex	CM3
150 mg	500 mg	orange/oblong, biconvex	CM2
150 mg	1,000 mg	reddish brown/oblong, biconvex	CM4

* Embossing appears on one side only of tablet.

4 CONTRAINDICATIONS

INVOKAMET/INVOKAMET XR is contraindicated in patients with:

- Moderate to severe renal impairment (eGFR below 45 mL/min/1.73 m²), end stage renal disease (ESRD) or patients on dialysis [see *Warnings and Precautions (5.1) and Use in Specific Populations (8.6)*].
- Acute or chronic metabolic acidosis, including diabetic ketoacidosis [see *Warnings and Precautions (5.4)*].
- Serious hypersensitivity reaction to canagliflozin or metformin, such as anaphylaxis or angioedema [see *Warnings and Precautions (5.10) and Adverse Reactions (6)*].

5 WARNINGS AND PRECAUTIONS

5.1 Lactic Acidosis

There have been post-marketing cases of metformin-associated lactic acidosis, including fatal cases. These cases had a subtle onset and were accompanied by nonspecific symptoms such as malaise, myalgias, abdominal pain, respiratory distress, or increased somnolence; however, hypothermia, hypotension and resistant bradyarrhythmias have occurred with severe acidosis. Metformin-associated lactic acidosis was characterized by elevated blood lactate concentrations (>5 mmol/Liter), anion gap acidosis (without evidence of ketonuria or ketonemia), and an increased lactate:pyruvate ratio; metformin plasma levels generally >5 mcg/mL. Metformin decreases liver uptake of lactate increasing lactate blood levels which may increase the risk of lactic acidosis, especially in patients at risk.

If metformin-associated lactic acidosis is suspected, general supportive measures should be instituted promptly in a hospital setting, along with immediate discontinuation of INVOKAMET/INVOKAMET XR. In INVOKAMET/INVOKAMET XR-treated patients with a diagnosis or strong suspicion of lactic acidosis, prompt hemodialysis is recommended to correct the acidosis and remove accumulated metformin (metformin hydrochloride is dialyzable, with a clearance of up to 170 mL/minute under good hemodynamic conditions). Hemodialysis has often resulted in reversal of symptoms and recovery.

Educate patients and their families about the symptoms of lactic acidosis and if these symptoms occur instruct them to discontinue INVOKAMET/INVOKAMET XR and report these symptoms to their healthcare provider.

For each of the known and possible risk factors for metformin-associated lactic acidosis, recommendations to reduce the risk of and manage metformin-associated lactic acidosis are provided below:

Renal Impairment: The postmarketing metformin-associated lactic acidosis cases primarily occurred in patients with significant renal impairment. The risk of metformin accumulation and metformin-associated lactic acidosis increases with the severity of renal impairment because metformin is substantially excreted by the kidney. Clinical recommendations based upon the patient's renal function include [see *Dosage and Administration (2.4), Clinical Pharmacology (12.3)*].

- Before initiating INVOKAMET/INVOKAMET XR, obtain an estimated glomerular filtration rate (eGFR).
- INVOKAMET/INVOKAMET XR is contraindicated in patients with an eGFR less than 45 mL/minute/1.73 m² [see *Contraindications (4)*].
- Obtain an eGFR at least annually in all patients taking INVOKAMET/INVOKAMET XR. In patients at increased risk for the development of renal impairment (e.g., the elderly), renal function should be assessed more frequently.

Drug Interactions: The concomitant use of INVOKAMET/INVOKAMET XR with specific drugs may increase the risk of metformin-associated lactic acidosis: those that impair renal function, result in significant hemodynamic change, interfere with acid-base balance or increase metformin accumulation (e.g. cationic drugs) [see *Drug Interactions (7)*]. Therefore, consider more frequent monitoring of patients.

Age 65 or Greater: The risk of metformin-associated lactic acidosis increases with the patient's age because elderly patients have a greater likelihood of having hepatic, renal, or cardiac impairment than younger patients. Assess renal function more frequently in elderly patients [see *Use in Specific Populations (8.5)*].

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Radiological Studies with Contrast: Administration of intravascular iodinated contrast agents in metformin-treated patients has led to an acute decrease in renal function and the occurrence of lactic acidosis. Stop INVOKAMET/INVOKAMET XR at the time of, or prior to, an iodinated contrast imaging procedure in patients with an eGFR between 45 and 60 mL/min/1.73 m²; in patients with a history of hepatic impairment, alcoholism, or heart failure; or in patients who will be administered intra-arterial iodinated contrast. Re-evaluate eGFR 48 hours after the imaging procedure, and restart INVOKAMET/INVOKAMET XR if renal function is stable.

Surgery and Other Procedures: Withholding of food and fluids during surgical or other procedures may increase the risk for volume depletion, hypotension and renal impairment.

INVOKAMET/INVOKAMET XR should be temporarily discontinued while patients have restricted food and fluid intake.

Hypoxic States: Several of the postmarketing cases of metformin-associated lactic acidosis occurred in the setting of acute congestive heart failure (particularly when accompanied by hypoperfusion and hypoxemia). Cardiovascular collapse (shock), acute myocardial infarction, sepsis, and other conditions associated with hypoxemia have been associated with lactic acidosis and may also cause pre-renal azotemia. When such events occur, discontinue INVOKAMET/INVOKAMET XR.

Excessive Alcohol Intake: Alcohol potentiates the effect of metformin on lactate metabolism and this may increase the risk of metformin-associated lactic acidosis. Warn patients against excessive alcohol intake while receiving INVOKAMET/INVOKAMET XR.

Hepatic Impairment: Patients with hepatic impairment have developed metformin-associated lactic acidosis. This may be due to impaired lactate clearance resulting in higher lactate blood levels. Therefore, avoid use of INVOKAMET/INVOKAMET XR in patients with clinical or laboratory evidence of hepatic disease.

5.2 Lower Limb Amputation

An approximately 2-fold increased risk of lower limb amputations associated with canagliflozin, a component of INVOKAMET/INVOKAMET XR, was observed in CANVAS and CANVAS-R, two randomized, placebo-controlled trials evaluating patients with type 2 diabetes who had either established cardiovascular disease or were at risk for cardiovascular disease. The risk of lower limb amputations was observed at both the 100 mg and 300 mg once daily dosage regimens. The amputation data for CANVAS and CANVAS-R are shown in Tables 2 and 3, respectively [see *Adverse Reactions* (6.1)].

Amputations of the toe and midfoot (99 out of 140 patients with amputations receiving canagliflozin in the two trials) were the most frequent; however, amputations involving the leg, below and above the knee, were also observed (41 out of 140 patients with amputations receiving canagliflozin in the two trials). Some patients had multiple amputations, some involving both lower limbs.

Lower limb infections, gangrene, and diabetic foot ulcers were the most common precipitating medical events leading to the need for an amputation. The risk of amputation was highest in patients with a baseline history of prior amputation, peripheral vascular disease, and neuropathy.

Before initiating INVOKAMET/INVOKAMET XR, consider factors in the patient history that may predispose to the need for amputations, such as a history of prior amputation, peripheral vascular disease, neuropathy and diabetic foot ulcers. Counsel patients about the importance of routine preventative foot care. Monitor patients receiving INVOKAMET/INVOKAMET XR for signs and symptoms of infection (including osteomyelitis), new pain or tenderness, sores or ulcers involving the lower limbs, and discontinue INVOKAMET/INVOKAMET XR if these complications occur.

5.3 Hypotension

Canagliflozin causes intravascular volume contraction. Symptomatic hypotension can occur after initiating INVOKAMET/INVOKAMET XR [see *Adverse Reactions* (6.1)] particularly in patients with impaired renal function (eGFR less than 60 mL/min/1.73 m²), elderly patients, patients on either diuretics or medications that interfere with the renin-angiotensin-aldosterone system (e.g., angiotensin-converting-enzyme [ACE] inhibitors, angiotensin receptor blockers [ARBs]), or patients with low systolic blood pressure. Before initiating INVOKAMET/INVOKAMET XR in patients with one or more of these characteristics who were not already on canagliflozin, volume status should be assessed and corrected. Monitor for signs and symptoms after initiating therapy.

5.4 Ketoacidosis

Reports of ketoacidosis, a serious life-threatening condition requiring urgent hospitalization have been identified in postmarketing surveillance in patients with type 1 and type 2 diabetes mellitus receiving sodium glucose co-transporter-2 (SGLT2) inhibitors, including canagliflozin. Fatal cases of ketoacidosis have been reported in patients taking canagliflozin. INVOKAMET/INVOKAMET XR is not indicated for the treatment of patients with type 1 diabetes mellitus [see *Indications and Usage* (1)].

Patients treated with INVOKAMET/INVOKAMET XR who present with signs and symptoms consistent with severe metabolic acidosis should be assessed for ketoacidosis regardless of presenting blood glucose levels, as ketoacidosis associated with INVOKAMET/INVOKAMET XR may be present even if blood

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

glucose levels are less than 250 mg/dL. If ketoacidosis is suspected, INVOKAMET/INVOKAMET XR should be discontinued, patient should be evaluated, and prompt treatment should be instituted. Treatment of ketoacidosis may require insulin, fluid and carbohydrate replacement.

In many of the postmarketing reports, and particularly in patients with type 1 diabetes, the presence of ketoacidosis was not immediately recognized and institution of treatment was delayed because presenting blood glucose levels were below those typically expected for diabetic ketoacidosis (often less than 250 mg/dL). Signs and symptoms at presentation were consistent with dehydration and severe metabolic acidosis and included nausea, vomiting, abdominal pain, generalized malaise, and shortness of breath. In some but not all cases, factors predisposing to ketoacidosis such as insulin dose reduction, acute febrile illness, reduced caloric intake due to illness or surgery, pancreatic disorders suggesting insulin deficiency (e.g., type 1 diabetes, history of pancreatitis or pancreatic surgery), and alcohol abuse were identified.

Before initiating INVOKAMET/INVOKAMET XR consider factors in the patient history that may predispose to ketoacidosis including pancreatic insulin deficiency from any cause, caloric restriction, and alcohol abuse. In patients treated with INVOKAMET/INVOKAMET XR consider monitoring for ketoacidosis and temporarily discontinuing INVOKAMET/INVOKAMET XR in clinical situations known to predispose to ketoacidosis (e.g., prolonged fasting due to acute illness or surgery).

5.5 Acute Kidney Injury

Canagliflozin causes intravascular volume contraction [see *Warnings and Precautions* (5.3)] and can cause acute kidney injury. There have been postmarketing reports of acute kidney injury, some requiring hospitalization and dialysis, in patients receiving canagliflozin; some reports involved patients younger than 65 years of age.

Before initiating INVOKAMET/INVOKAMET XR, consider factors that may predispose patients to acute kidney injury including hypovolemia, chronic renal insufficiency, congestive heart failure, and concomitant medications (diuretics, ACE inhibitors, ARBs, NSAIDs). Consider temporarily discontinuing INVOKAMET/INVOKAMET XR in any setting of reduced oral intake (such as acute illness or fasting) or fluid losses (such as gastrointestinal illness or excessive heat exposure); monitor patients for signs and symptoms of acute kidney injury. If acute kidney injury occurs, discontinue INVOKAMET/INVOKAMET XR promptly and institute treatment.

Initiation of canagliflozin may increase serum creatinine and decrease eGFR. Patients with hypovolemia may be more susceptible to these changes. [see *Adverse Reactions* (6.1)]. Renal function should be evaluated prior to initiation of INVOKAMET/INVOKAMET XR and monitored periodically thereafter. Dosage adjustment and more frequent renal function monitoring are recommended in patients with an eGFR below 60 mL/min/1.73 m². INVOKAMET/INVOKAMET XR is contraindicated in patients with an eGFR below 45 mL/min/1.73 m² [see *Dosage and Administration* (2.4), *Contraindications* (4), *Warnings and Precautions* (5.1) and *Use in Specific Populations* (8.6)].

5.6 Urrosepsis and Pyelonephritis

There have been postmarketing reports of serious urinary tract infections including urrosepsis and pyelonephritis requiring hospitalization in patients receiving SGLT2 inhibitors, including canagliflozin. Treatment with SGLT2 inhibitors increases the risk for urinary tract infections. Evaluate patients for signs and symptoms of urinary tract infections and treat promptly, if indicated [see *Adverse Reactions* (6)].

5.7 Hypoglycemia with Concomitant Use of Sulfonylurea or Insulin

Insulin and insulin secretagogues are known to cause hypoglycemia. INVOKAMET/INVOKAMET XR may increase the risk of hypoglycemia when combined with insulin or an insulin secretagogue [see *Adverse Reactions* (6.1)]. Therefore, a lower dose of insulin or insulin secretagogue may be required to minimize the risk of hypoglycemia when used in combination with INVOKAMET/INVOKAMET XR.

5.8 Necrotizing Fasciitis of the Perineum (Fournier's Gangrene)

Reports of necrotizing fasciitis of the perineum (Fournier's gangrene), a rare but serious and life-threatening necrotizing infection requiring urgent surgical intervention, have been identified in postmarketing surveillance in patients with diabetes mellitus receiving SGLT2 inhibitors, including canagliflozin. Cases have been reported in both females and males. Serious outcomes have included hospitalization, multiple surgeries, and death.

Patients treated with INVOKAMET/INVOKAMET XR presenting with pain or tenderness, erythema, or swelling in the genital or perineal area, along with fever or malaise, should be assessed for necrotizing fasciitis. If suspected, start treatment immediately with broad-spectrum antibiotics and, if necessary, surgical debridement. Discontinue INVOKAMET/INVOKAMET XR, closely monitor blood glucose levels, and provide appropriate alternative therapy for glycemic control.

5.9 Genital Mycotic Infections

Canagliflozin increases the risk of genital mycotic infections. Patients with a history of genital mycotic infections and uncircumcised males were more likely to develop genital mycotic infections [see *Adverse Reactions* (6.1)]. Monitor and treat appropriately.

5.10 Hypersensitivity Reactions

Hypersensitivity reactions, including angioedema and anaphylaxis, have been reported with canagliflozin. These reactions generally occurred within hours to days after initiating canagliflozin. If hypersensitivity reactions occur, discontinue use of INVOKAMET/INVOKAMET XR; treat and monitor until signs and symptoms resolve [see *Contraindications (4) and Adverse Reactions (6.1, 6.2)*].

5.11 Bone Fracture

An increased risk of bone fracture, occurring as early as 12 weeks after treatment initiation, was observed in patients using canagliflozin in the CANVAS trial [see *Clinical Studies (14.2)*]. Consider factors that contribute to fracture risk prior to initiating INVOKAMET/INVOKAMET XR [see *Adverse Reactions (6.1)*].

5.12 Vitamin B₁₂ Levels

In metformin HCl clinical trials of 29-week duration, a decrease to subnormal levels of previously normal serum vitamin B₁₂ levels was observed in approximately 7% of patients. Such decrease, possibly due to interference with B₁₂ absorption from the B₁₂-intrinsic factor complex, may be associated with anemia but appears to be rapidly reversible with discontinuation of metformin or vitamin B₁₂ supplementation. Certain individuals (those with inadequate vitamin B₁₂ or calcium intake or absorption) appear to be predisposed to developing subnormal vitamin B₁₂ levels. Measure hematologic parameters on an annual basis and vitamin B₁₂ at 2- to 3-year intervals in patients on INVOKAMET/INVOKAMET XR and manage any abnormalities [see *Adverse Reactions (6.1)*].

5.13 Increases in Low-Density Lipoprotein (LDL-C)

Dose-related increases in LDL-C occur with canagliflozin [see *Adverse Reactions (6.1)*]. Monitor LDL-C and treat if appropriate after initiating INVOKAMET/INVOKAMET XR.

6 ADVERSE REACTIONS

The following important adverse reactions are also discussed elsewhere in the labeling:

- Lactic Acidosis [see *Boxed Warning and Warnings and Precautions (5.1, 5.5)*]
- Lower Limb Amputation [see *Boxed Warning and Warnings and Precautions (5.2)*]
- Hypotension [see *Warnings and Precautions (5.3)*]
- Ketoacidosis [see *Warnings and Precautions (5.4)*]
- Acute Kidney Injury [see *Warnings and Precautions (5.5)*]
- Urosepsis and Pyelonephritis [see *Warnings and Precautions (5.6)*]
- Hypoglycemia with Concomitant Use of Sulfonylurea or Insulin [see *Warnings and Precautions (5.7)*]
- Necrotizing Fasciitis of the Perineum (Fournier's gangrene) [see *Warnings and Precautions (5.8)*]
- Genital Mycotic Infections [see *Warnings and Precautions (5.9)*]
- Hypersensitivity Reactions [see *Warnings and Precautions (5.10)*]
- Bone Fracture [see *Warnings and Precautions (5.11)*]
- Vitamin B₁₂ Deficiency [see *Warnings and Precautions (5.12)*]
- Increases in Low-Density Lipoprotein (LDL-C) [see *Warnings and Precautions (5.13)*]

6.1 Clinical Studies Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to the rates in the clinical trials of another drug and may not reflect the rates observed in clinical practice.

Pool of Placebo-Controlled Trials

Canagliflozin

The data in Table 1 is derived from four 26-week placebo-controlled trials where canagliflozin was used as monotherapy in one trial and as add-on therapy in three trials. These data reflect exposure of 1,667 patients to canagliflozin and a mean duration of exposure to canagliflozin of 24 weeks with 1,275 patients exposed to a combination of canagliflozin and metformin. Patients received canagliflozin 100 mg (N=833), canagliflozin 300 mg (N=834) or placebo (N=646) once daily. The mean daily dose of metformin HCl was 2,138 mg (SD 337.3) for the 1,275 patients in the three placebo-controlled metformin add-on trials. The mean age of the population was 56 years and 2% were older than 75 years of age. Fifty percent (50%) of the population was male and 72% were Caucasian, 12% were Asian, and 5% were Black or African American. At baseline the population had diabetes for an average of 7.3 years, had a mean HbA_{1c} of 8.0% and 20% had established microvascular complications of diabetes. Baseline renal function was normal or mildly impaired (mean eGFR 88 mL/min/1.73 m²).

Table 1 shows common adverse reactions associated with the use of canagliflozin. These adverse reactions were not present at baseline, occurred more commonly on canagliflozin than on placebo, and occurred in at least 2% of patients treated with either canagliflozin 100 mg or canagliflozin 300 mg.

Table 1: Adverse Reactions From Pool of Four 26-Week Placebo-Controlled Studies Reported in ≥ 2% of Canagliflozin-Treated Patients*

Adverse Reaction	Placebo N=646	Canagliflozin 100 mg N=833	Canagliflozin 300 mg N=834
Urinary tract infections [‡]	3.8%	5.9%	4.4%
Increased urination [§]	0.7%	5.1%	4.6%
Thirst [¶]	0.1%	2.8%	2.4%
Constipation	0.9%	1.8%	2.4%
Nausea	1.6%	2.1%	2.3%
	N=312	N=425	N=430
Female genital mycotic infections [†]	2.8%	10.6%	11.6%
Vulvovaginal pruritus	0.0%	1.6%	3.2%
	N=334	N=408	N=404
Male genital mycotic infections ^{¶¶}	0.7%	4.2%	3.8%

* The four placebo-controlled trials included one monotherapy trial and three add-on combination trials with metformin, metformin and sulfonylurea, or metformin and pioglitazone.

[†] Female genital mycotic infections include the following adverse reactions: Vulvovaginal candidiasis, Vulvovaginal mycotic infection, Vulvovaginitis, Vaginal infection, Vulvitis, and Genital infection fungal.

[‡] Urinary tract infections include the following adverse reactions: Urinary tract infection, Cystitis, Kidney infection, and Urosepsis.

[§] Increased urination includes the following adverse reactions: Polyuria, Pollakiuria, Urine output increased, Micturition urgency, and Nocturia.

[¶] Male genital mycotic infections include the following adverse reactions: Balanitis or Balanoposthitis, Balanitis candida, and Genital infection fungal.

^{¶¶} Thirst includes the following adverse reactions: Thirst, Dry mouth, and Polydipsia. Note: Percentages were weighted by studies. Study weights were proportional to the harmonic mean of the three treatment sample sizes.

Abdominal pain was also more commonly reported in patients taking canagliflozin 100 mg (1.8%), 300 mg (1.7%) than in patients taking placebo (0.8%).

Canagliflozin and Metformin

The incidence and type of adverse reactions in the three 26-week placebo-controlled metformin add-on trials, representing a majority of data from the four 26-week placebo-controlled trials, was similar to the adverse reactions described in Table 1. There were no additional adverse reactions identified in the pooling of these three placebo-controlled trials that included metformin relative to the four placebo-controlled trials.

In a trial with canagliflozin as initial combination therapy with metformin [see *Clinical Studies (14.1)*], an increased incidence of diarrhea was observed in the canagliflozin and metformin combination groups (4.2%) compared to canagliflozin or metformin monotherapy groups (1.7%).

Pool of Placebo- and Active-Controlled Trials - Canagliflozin

The occurrence of adverse reactions for canagliflozin was evaluated in patients participating in placebo- and active-controlled trials and in an integrated analysis of two cardiovascular trials.

The types and frequency of common adverse reactions observed in the pool of eight clinical trials (which reflect an exposure of 6,177 patients to canagliflozin) were consistent with those listed in Table 1. Percentages were weighted by studies. Study weights were proportional to the harmonic mean of the three treatment sample sizes. In this pool, canagliflozin was also associated with the adverse reactions of fatigue (1.8%, 2.2%, and 2.0% with comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively) and loss of strength or energy (i.e., asthenia) (0.6%, 0.7%, and 1.1% with comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively).

In the pool of eight clinical trials, the incidence rate of pancreatitis (acute or chronic) was 0.1%, 0.2%, and 0.1% receiving comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

In the pool of eight clinical trials, hypersensitivity-related adverse reactions (including erythema, rash, pruritus, urticaria, and angioedema) was 3.0%, 3.8%, and 4.2% of patients receiving comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Five patients experienced serious adverse reactions of hypersensitivity with canagliflozin, which included 4 patients with urticaria and 1 patient with a diffuse rash and urticaria occurring within hours of exposure to canagliflozin. Among these patients, 2 patients discontinued canagliflozin. One patient with urticaria had recurrence when canagliflozin was re-initiated.

Photosensitivity-related adverse reactions (including photosensitivity reaction, polymorphic light eruption, and sunburn) occurred in 0.1%, 0.2%, and 0.2% of patients receiving comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Other adverse reactions occurring more frequently on canagliflozin than on comparator were:

Lower Limb Amputation

An approximately 2-fold increased risk of lower limb amputations associated with canagliflozin was observed in CANVAS and CANVAS-R, two randomized, placebo-controlled trials evaluating patients with type 2 diabetes who had either established cardiovascular disease or were at risk for cardiovascular disease. Patients in CANVAS and CANVAS-R were followed for an average of 5.7 and 2.1 years, respectively [see *Clinical Studies (14.2)*]. The amputation data for CANVAS and CANVAS-R are shown in Tables 2 and 3, respectively [see *Warnings and Precautions (5.2)*].

Table 2: CANVAS Amputations

	Placebo N=1441	Canagliflozin 100 mg N=1445	Canagliflozin 300 mg N=1441	Canagliflozin (Pooled) N=2886
Patients with an amputation, n (%)	22 (1.5)	50 (3.5)	45 (3.1)	95 (3.3)
Total amputations	33	83	79	162
Amputation incidence rate (per 1000 patient-years)	2.8	6.2	5.5	5.9
Hazard Ratio (95% CI)	--	2.24 (1.36, 3.69)	2.01 (1.20, 3.34)	2.12 (1.34, 3.38)

Note: Incidence is based on the number of patients with at least one amputation, and not the total number of amputation events. A patient's follow-up is calculated from Day 1 to the first amputation event date. Some patients had more than one amputation.

Table 3: CANVAS-R Amputations

	Placebo N=2903	Canagliflozin 100 mg (with up-titration to 300 mg) N=2904
Patients with an amputation, n (%)	25 (0.9)	45 (1.5)
Total amputations	36	59
Amputation incidence rate (per 1000 patient-years)	4.2	7.5
Hazard Ratio (95% CI)	--	1.80 (1.10, 2.93)

Note: Incidence is based on the number of patients with at least one amputation, and not the total number of amputation events. A patient's follow-up is calculated from Day 1 to the first amputation event date. Some patients had more than one amputation.

Renal Cell Carcinoma

In the CANVAS trial (mean duration of follow-up of 5.7 years) [see *Clinical Studies (14.2)*], the incidence of renal cell carcinoma was 0.15% (2/1331) and 0.29% (8/2716) for placebo and canagliflozin, respectively, excluding patients with less than 6 months of follow-up, less than 90 days of treatment, or a history of renal cell carcinoma. A causal relationship to canagliflozin could not be established due to the limited number of cases.

Volume Depletion-Related Adverse Reactions

Canagliflozin results in an osmotic diuresis, which may lead to reductions in intravascular volume. In clinical trials, treatment with canagliflozin was associated with a dose-dependent increase in the incidence of volume depletion-related adverse reactions (e.g., hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration). An increased incidence was observed in patients on the 300 mg dose. The three factors associated with the largest increase in volume depletion-related adverse reactions were the use of loop diuretics, moderate renal impairment (eGFR 30 to less than 60 mL/min/1.73 m²), and age 75 years and older (Table 4) [see *Dosage and Administration (2.1)*, *Warnings and Precautions (5.3)*, and *Use in Specific Populations (8.5 and 8.6)*].

Table 4: Patients With at Least One Volume Depletion-Related Adverse Reaction (Pooled Results from 8 Clinical Trials)

Baseline Characteristic	Comparator Group* %	Canagliflozin 100 mg %	Canagliflozin 300 mg %
Overall population	1.5%	2.3%	3.4%
75 years of age and older [†]	2.6%	4.9%	8.7%
eGFR less than 60 mL/min/1.73 m ^{2†}	2.5%	4.7%	8.1%
Use of loop diuretic [†]	4.7%	3.2%	8.8%

* Includes placebo and active-comparator groups

[†] Patients could have more than 1 of the listed risk factors

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Falls

In a pool of nine clinical trials with mean duration of exposure to canagliflozin of 85 weeks, the proportion of patients who experienced falls was 1.3%, 1.5%, and 2.1% with comparator, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. The higher risk of falls for patients treated with canagliflozin was observed within the first few weeks of treatment.

Impairment in Renal Function

Initiation of canagliflozin is associated with a dose-dependent increase in serum creatinine and a concomitant fall in estimated GFR (Table 5) [see *Warnings and Precautions (5.5)*]. The effect on eGFR was observed to reverse after treatment discontinuation suggesting acute hemodynamic changes may play a role in the renal function changes observed with canagliflozin.

Table 5: Changes in Serum Creatinine and eGFR Associated with Canagliflozin in the Pool of Four Placebo-Controlled Trials and Moderate Renal Impairment Trial

			Placebo N=646	Canagliflozin 100 mg N=833	Canagliflozin 300 mg N=834
Pool of Four Placebo-Controlled Trials	Baseline	Creatinine (mg/dL)	0.84	0.82	0.82
		eGFR (mL/min/1.73 m ²)	87.0	88.3	88.8
	Week 6 Change	Creatinine (mg/dL)	0.01	0.03	0.05
		eGFR (mL/min/1.73 m ²)	-1.6	-3.8	-5.0
	End of Treatment Change*	Creatinine (mg/dL)	0.01	0.02	0.03
		eGFR (mL/min/1.73 m ²)	-1.6	-2.3	-3.4
			Placebo N=90	Canagliflozin 100 mg N=90	Canagliflozin 300 mg N=89
Moderate Renal Impairment Trial	Baseline	Creatinine (mg/dL)	1.61	1.62	1.63
		eGFR (mL/min/1.73 m ²)	40.1	39.7	38.5
	Week 3 Change	Creatinine (mg/dL)	0.03	0.18	0.28
		eGFR (mL/min/1.73 m ²)	-0.7	-4.6	-6.2
	End of Treatment Change*	Creatinine (mg/dL)	0.07	0.16	0.18
		eGFR (mL/min/1.73 m ²)	-1.5	-3.6	-4.0

* Week 26 in mITT LOCF population

In the pool of four placebo-controlled trials where patients had normal or mildly impaired baseline renal function, the proportion of patients who experienced at least one event of significant renal function decline, defined as an eGFR below 80 mL/min/1.73 m² and 30% lower than baseline, was 2.1% with placebo, 2.0% with canagliflozin 100 mg, and 4.1% with canagliflozin 300 mg. At the end of treatment, 0.5% with placebo, 0.7% with canagliflozin 100 mg, and 1.4% with canagliflozin 300 mg had a significant renal function decline.

Patients with moderate renal impairment at baseline experience larger mean changes in eGFR relative to patients with normal or mildly impaired renal function. In a trial in patients with moderate renal impairment with a baseline eGFR of 30 to less than 50 mL/min/1.73 m² (mean baseline eGFR 39 mL/min/1.73 m²) [see *Clinical Studies (14.1)*], the proportion of patients who experienced at least one event of significant renal function decline, defined as an eGFR 30% lower than baseline, was 6.9%, 18%, and 22.5% with placebo, canagliflozin 100 mg, canagliflozin 300 mg, respectively. At the end of treatment, 4.6%, 3.4%, and 2.2% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively, had a significant renal function decline.

Genital Mycotic Infections

In the pool of four placebo-controlled clinical trials, female genital mycotic infections (e.g., vulvovaginal mycotic infection, vulvovaginal candidiasis, and vulvovaginitis) occurred in 2.8%, 10.6%, and 11.6% of females treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Patients with a history of genital mycotic infections were more likely to develop genital mycotic infections on canagliflozin. Female patients who developed genital mycotic infections on canagliflozin were more likely to experience recurrence and require treatment with oral or topical antifungal agents and anti-microbial agents. In females, discontinuation due to genital mycotic infections occurred in 0% and 0.7% of patients treated with placebo and canagliflozin, respectively [see *Warnings and Precautions (5.9)*].

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

In the pool of four placebo-controlled clinical trials, male genital mycotic infections (e.g., candidal balanitis, balanoposthitis) occurred in 0.7%, 4.2%, and 3.8% of males treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Male genital mycotic infections occurred more commonly in uncircumcised males and in males with a prior history of balanitis or balanoposthitis. Male patients who developed genital mycotic infections on canagliflozin were more likely to experience recurrent infections (22% on canagliflozin versus none on placebo), and require treatment with oral or topical antifungal agents and anti-microbial agents than patients on comparators. In males, discontinuations due to genital mycotic infections occurred in 0% and 0.5% of patients treated with placebo and canagliflozin, respectively.

In the pooled analysis of 8 controlled trials, phimosis was reported in 0.3% of uncircumcised male patients treated with canagliflozin and 0.2% required circumcision to treat the phimosis [see *Warnings and Precautions (5.9)*].

Hypoglycemia

In canagliflozin clinical trials, hypoglycemia was defined as any event regardless of symptoms, where biochemical hypoglycemia was documented (any glucose value below or equal to 70 mg/dL). Severe hypoglycemia was defined as an event consistent with hypoglycemia where the patient required the assistance of another person to recover, lost consciousness, or experienced a seizure (regardless of whether biochemical documentation of a low glucose value was obtained). In individual clinical trials [see *Clinical Studies (14.1)*], episodes of hypoglycemia occurred at a higher rate when canagliflozin was co-administered with insulin or sulfonylureas (Table 6) [see *Warnings and Precautions (5.7)*].

Table 6: Incidence of Hypoglycemia* in Controlled Clinical Studies

Monotherapy (26 weeks)	Placebo (N=192)	Canagliflozin 100 mg (N=195)	Canagliflozin 300 mg (N=197)
Overall [N (%)]	5 (2.6)	7 (3.6)	6 (3.0)
In Combination with Metformin (26 weeks)	Placebo + Metformin (N=183)	Canagliflozin 100 mg + Metformin (N=368)	Canagliflozin 300 mg + Metformin (N=367)
Overall [N (%)]	3 (1.6)	16 (4.3)	17 (4.6)
Severe [N (%)] [†]	0 (0)	1 (0.3)	1 (0.3)
In Combination with Metformin (18 weeks)[‡]	Placebo (N=93)	Canagliflozin 100 mg (N=93)	Canagliflozin 300 mg (N=93)
Overall [N (%)]	3 (3.2)	4 (4.3)	3 (3.2)
In Combination with Metformin + Sulfonylurea (26 weeks)	Placebo + Metformin + Sulfonylurea (N=156)	Canagliflozin 100 mg + Metformin + Sulfonylurea (N=157)	Canagliflozin 300 mg + Metformin + Sulfonylurea (N=156)
Overall [N (%)]	24 (15.4)	43 (27.4)	47 (30.1)
Severe [N (%)] [†]	1 (0.6)	1 (0.6)	0
In Combination with Metformin + Pioglitazone (26 weeks)	Placebo + Metformin + Pioglitazone (N=115)	Canagliflozin 100 mg + Metformin + Pioglitazone (N=113)	Canagliflozin 300 mg + Metformin + Pioglitazone (N=114)
Overall [N (%)]	3 (2.6)	3 (2.7)	6 (5.3)
In Combination with Insulin (18 weeks)	Placebo (N=565)	Canagliflozin 100 mg (N=566)	Canagliflozin 300 mg (N=587)
Overall [N (%)]	208 (36.8)	279 (49.3)	285 (48.6)
Severe [N (%)] [†]	14 (2.5)	10 (1.8)	16 (2.7)
In Combination with Insulin and Metformin (18 weeks)[§]	Placebo (N=145)	Canagliflozin 100 mg (N=139)	Canagliflozin 300 mg (N=148)
Overall [N (%)]	66 (45.5)	58 (41.7)	70 (47.3)
Severe [N (%)] [†]	4 (2.8)	1 (0.7)	3 (2.0)

* Number of patients experiencing at least one event of hypoglycemia based on either biochemically documented episodes or severe hypoglycemic events in the intent-to-treat population

[†] Severe episodes of hypoglycemia were defined as those where the patient required the assistance of another person to recover, lost consciousness, or experienced a seizure (regardless of whether biochemical documentation of a low glucose value was obtained)

[‡] Phase 2 clinical study with twice daily dosing (50 mg or 150 mg twice daily in combination with metformin)

[§] Subgroup of patients (N=287) from insulin substudy on canagliflozin in combination with metformin and insulin (with or without other antiglycemic agents)

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Bone Fracture

In the CANVAS trial [see *Clinical Studies (14.2)*], the incidence rates of all adjudicated bone fracture were 1.09, 1.59, and 1.79 events per 100 patient-years of follow-up to placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. The fracture imbalance was observed within the first 26 weeks of therapy and remained through the end of the trial. Fractures were more likely to be low trauma (e.g., fall from no more than standing height), and affect the distal portion of upper and lower extremities [see *Warnings and Precautions (5.11)*].

Metformin

The most common adverse reactions (5% or greater incidence) due to initiation of metformin are diarrhea, nausea, vomiting, flatulence, asthenia, indigestion, abdominal discomfort, and headache.

Long-term treatment with metformin has been associated with a decrease in vitamin B₁₂, which may result in clinically significant vitamin B₁₂ deficiency (e.g., megaloblastic anemia) [see *Warnings and Precautions (5.12)*].

Laboratory and Imaging Tests

Increases in Serum Potassium

In a pooled population of patients (N=723) with moderate renal impairment (eGFR 45 to less than 60 mL/min/1.73 m²), increases in serum potassium to greater than 5.4 mEq/L and 15% above occurred in 5.3%, 5.0%, and 8.8% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Severe elevations (greater than or equal to 6.5 mEq/L) occurred in 0.4% of patients treated with placebo, no patients treated with canagliflozin 100 mg, and 1.3% of patients treated with canagliflozin 300 mg.

In these patients, increases in potassium were more commonly seen in those with elevated potassium at baseline. Among patients with moderate renal impairment, approximately 84% were taking medications that interfere with potassium excretion, such as potassium-sparing diuretics, angiotensin-converting-enzyme inhibitors, and angiotensin-receptor blockers [see *Warnings and Precautions (5.3) and Use in Specific Populations (8.6)*].

Increases in Serum Magnesium

Dose-related increases in serum magnesium were observed early after initiation of canagliflozin (within 6 weeks) and remained elevated throughout treatment. In the pool of four placebo-controlled trials, the mean percent change in serum magnesium levels was 8.1% and 9.3% with canagliflozin 100 mg and canagliflozin 300 mg, respectively, compared to -0.6% with placebo. In a trial of patients with moderate renal impairment, serum magnesium levels increased by 0.2%, 9.2%, and 14.8% with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

Increases in Serum Phosphate

Dose-related increases in serum phosphate levels were observed with canagliflozin. In the pool of four placebo-controlled trials, the mean percent change in serum phosphate levels were 3.6% and 5.1% with canagliflozin 100 mg and canagliflozin 300 mg, respectively, compared to 1.5% with placebo. In a trial of patients with moderate renal impairment, the mean serum phosphate levels increased by 1.2%, 5.0%, and 9.3% with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively.

Increases in Low-Density Lipoprotein Cholesterol (LDL-C) and non-High-Density Lipoprotein Cholesterol (non-HDL-C)

In the pool of four placebo-controlled trials, dose-related increases in LDL-C with canagliflozin were observed. Mean changes (percent changes) from baseline in LDL-C relative to placebo were 4.4 mg/dL (4.5%) and 8.2 mg/dL (8.0%) with canagliflozin 100 mg and canagliflozin 300 mg, respectively. The mean baseline LDL-C levels were 104 to 110 mg/dL across treatment groups [see *Warnings and Precautions (5.13)*].

Dose-related increases in non-HDL-C with canagliflozin were observed. Mean changes (percent changes) from baseline in non-HDL-C relative to placebo were 2.1 mg/dL (1.5%) and 5.1 mg/dL (3.6%) with canagliflozin 100 mg and 300 mg, respectively. The mean baseline non-HDL-C levels were 140 to 147 mg/dL across treatment groups.

Increases in Hemoglobin

In the pool of four placebo-controlled trials, mean changes (percent changes) from baseline in hemoglobin were -0.18 g/dL (-1.1%) with placebo, 0.47 g/dL (3.5%) with canagliflozin 100 mg, and 0.51 g/dL (3.8%) with canagliflozin 300 mg. The mean baseline hemoglobin value was approximately 14.1 g/dL across treatment groups. At the end of treatment, 0.8%, 4.0%, and 2.7% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively, had hemoglobin above the upper limit of normal.

Decreases in Bone Mineral Density

Bone mineral density (BMD) was measured by dual-energy X-ray absorptiometry in a clinical trial of 714 older adults (mean age 64 years). At 2 years, patients randomized to canagliflozin 100 mg and canagliflozin 300 mg had placebo-corrected declines in BMD at the total hip of 0.9% and 1.2%, respectively, and at the lumbar spine of 0.3% and 0.7%, respectively. Additionally, placebo-adjusted BMD declines were 0.1% at the femoral neck for both canagliflozin doses and 0.4% at the distal forearm for patients randomized to canagliflozin 300 mg. The placebo-adjusted change at the distal forearm for patients randomized to canagliflozin 100 mg was 0%.

6.2 Postmarketing Experience

Additional adverse reactions have been identified during post-approval use of canagliflozin. Because these reactions are reported voluntarily from a population of uncertain size, it is generally not possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Canagliflozin

Ketoacidosis

Acute Kidney Injury

Anaphylaxis, Angioedema

Urosepsis and Pyelonephritis

Necrotizing Fasciitis of the Perineum (Fournier's gangrene)

Metformin

Cholestatic, hepatocellular, and mixed hepatocellular liver injury

7 DRUG INTERACTIONS

7.1 Drug Interactions with Metformin

Carbonic Anhydrase Inhibitors

Topiramate or other carbonic anhydrase inhibitors (e.g., zonisamide, acetazolamide or dichlorphenamide) frequently causes a decrease in serum bicarbonate and induce non-anion gap, hyperchloremic metabolic acidosis. Concomitant use of these drugs with INVOKAMET/INVOKAMET XR may increase the risk for lactic acidosis. Consider more frequent monitoring of these patients.

Drugs That Reduce Metformin Clearance

Concomitant use of drugs that interfere with common renal tubular transport systems involved in the renal elimination of metformin (e.g., organic cationic transporter-2 [OCT2] / multidrug and toxin extrusion [MATE] inhibitors such as ranolazine, vandetanib, dolutegravir, and cimetidine) could increase systemic exposure to metformin and may increase the risk for lactic acidosis [see *Clinical Pharmacology (12.3)*]. Consider the benefits and risks of concomitant use.

Alcohol

Alcohol is known to potentiate the effect of metformin on lactate metabolism. Warn patients against excessive alcohol intake while receiving INVOKAMET/INVOKAMET XR.

Drugs Affecting Glycemic Control

Certain drugs tend to produce hyperglycemia and may lead to loss of glycemic control. These drugs include the thiazides and other diuretics, corticosteroids, phenothiazines, thyroid products, estrogens, oral contraceptives, phenytoin, nicotinic acid, sympathomimetics, calcium channel blockers, and isoniazid. When such drugs are administered to a patient receiving INVOKAMET/INVOKAMET XR, monitor for loss of blood glucose control. When such drugs are withdrawn from a patient receiving INVOKAMET/INVOKAMET XR, monitor for hypoglycemia.

7.2 Drug Interactions with Canagliflozin

UGT Enzyme Inducers

Rifampin: Rifampin lowered canagliflozin exposure which may reduce the efficacy of INVOKAMET/INVOKAMET XR. If an inducer of UGTs (e.g., rifampin, phenytoin, phenobarbital, ritonavir) must be co-administered with INVOKAMET/INVOKAMET XR, consider increasing the total daily dose of canagliflozin to 300 mg if patients are currently tolerating INVOKAMET/INVOKAMET XR with a total daily dose of canagliflozin 100 mg, have an eGFR greater than 60 mL/min/1.73 m², and require additional glycemic control. Consider other antihyperglycemic therapy in patients with an eGFR of 45 to less than 60 mL/min/1.73 m² receiving concurrent therapy with a UGT inducer and require additional glycemic control [see *Dosage and Administration (2.5) and Clinical Pharmacology (12.3)*].

Digoxin

Canagliflozin increased digoxin exposure. Digoxin, as a cationic drug, also has the potential to compete with metformin for common renal tubular transport systems [see *Clinical Pharmacology (12.3)*]. Monitor patients taking INVOKAMET/INVOKAMET XR with concomitant digoxin for a need to adjust dose of either drug.

Drug/Laboratory Test Interference

Positive Urine Glucose Test

Monitoring glycemic control with urine glucose tests is not recommended in patients taking SGLT2 inhibitors as SGLT2 inhibitors increase urinary glucose excretion and will lead to positive urine glucose tests. Use alternative methods to monitor glycemic control.

Interference with 1,5-anhydroglucitol (1,5-AG) Assay

Monitoring glycemic control with 1,5-AG assay is not recommended as measurements of 1,5-AG are unreliable in assessing glycemic control in patients taking SGLT2 inhibitors. Use alternative methods to monitor glycemic control.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

Based on animal data showing adverse renal effects, INVOKAMET/INVOKAMET XR is not recommended during the second and third trimesters of pregnancy.

Limited data with INVOKAMET, INVOKAMET XR or canagliflozin in pregnant women are not sufficient to determine a drug-associated risk for major birth defects or miscarriage. Published studies with metformin use during pregnancy have not reported a clear association with metformin and major birth defect or miscarriage risk [see *Data*]. There are risks to the mother and fetus associated with poorly controlled diabetes in pregnancy [see *Clinical Considerations*].

In animal studies, adverse renal pelvic and tubule dilatations that were not reversible were observed in rats when canagliflozin was administered at an exposure 0.5-times the 300 mg clinical dose, based on AUC during a period of renal development corresponding to the late second and third trimesters of human pregnancy. No adverse developmental effects were observed when metformin was administered to pregnant Sprague Dawley rats and rabbits during the period of organogenesis at doses up to 2- and 6-times, respectively, a 2000 mg clinical dose, based on body surface area [see *Data*].

The estimated background risk of major birth defects is 6-10% in women with pre-gestational diabetes with an HbA_{1c} >7 and has been reported to be as high as 20-25% in women with a HbA_{1c} >10. The estimated background risk of miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Clinical Considerations

Disease-associated maternal and/or embryo/fetal risk

Poorly controlled diabetes in pregnancy increases the maternal risk for diabetic ketoacidosis, pre-eclampsia, spontaneous abortions, preterm delivery, stillbirth and delivery complications. Poorly controlled diabetes increases the fetal risk for major birth defects, stillbirth, and macrosomia related morbidity.

Data

Human Data

Published data from post-marketing studies have not reported a clear association with metformin and major birth defects, miscarriage, or adverse maternal or fetal outcomes when metformin was used during pregnancy. However, these studies cannot definitely establish the absence of any metformin-associated risk because of methodological limitations, including small sample size and inconsistent comparator groups.

Animal Data

Canagliflozin

Canagliflozin dosed directly to juvenile rats from postnatal day (PND) 21 until PND 90 at doses of 4, 20, 65, or 100 mg/kg increased kidney weights and dose dependently increased the incidence and severity of renal pelvic and tubular dilatation at all doses tested. Exposure at the lowest dose was greater than or equal to 0.5-times the 300 mg clinical dose, based on AUC. These outcomes occurred with drug exposure during periods of renal development in rats that correspond to the late second and third trimester of human renal development. The renal pelvic dilatations observed in juvenile animals did not fully reverse within a 1 month recovery period.

In embryo-fetal development studies in rats and rabbits, canagliflozin was administered for intervals coinciding with the first trimester period of organogenesis in humans. No developmental toxicities independent of maternal toxicity were observed when canagliflozin was administered at doses up to 100 mg/kg in pregnant rats and 160 mg/kg in pregnant rabbits during embryonic organogenesis or during a study in which maternal rats were dosed from gestation day (GD) 6 through PND 21, yielding exposures up to approximately 19-times the 300 mg clinical dose, based on AUC.

Metformin Hydrochloride

Metformin hydrochloride did not cause adverse developmental effects when administered to pregnant Sprague Dawley rats and rabbits up to 600 mg/kg/day during the period of organogenesis. This represents an exposure of about 2- and 6-times a 2000 mg clinical dose based on body surface area (mg/m²) for rats and rabbits, respectively.

Canagliflozin and Metformin

No adverse developmental effects were observed when canagliflozin and metformin were co-administered to pregnant rats during the period of organogenesis at exposures up to 11 and 13 times, respectively, the 300 mg and 2000 mg clinical doses of canagliflozin and metformin based on AUC.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

8.2 Lactation

Risk Summary

There is no information regarding the presence of INVOKAMET, INVOKAMET XR or canagliflozin in human milk, the effects on the breastfed infant, or the effects on milk production. Limited published studies report that metformin is present in human milk [see Data]. However, there is insufficient information on the effects of metformin on the breastfed infant and no available information on the effects of metformin on milk production. Canagliflozin is present in the milk of lactating rats [see Data]. Since human kidney maturation occurs *in utero* and during the first 2 years of life when lactational exposure may occur, there may be risk to the developing human kidney.

Because of the potential for serious adverse reactions in a breastfed infant, advise women that use of INVOKAMET/INVOKAMET XR is not recommended while breastfeeding.

Data

Human Data

Published clinical lactation studies report that metformin is present in human milk which resulted in infant doses approximately 0.11% to 1% of the maternal weight-adjusted dosage and a milk/plasma ratio ranging between 0.13 and 1. However, the studies were not designed to definitely establish the risk of use of metformin during lactation because of small sample size and limited adverse event data collected in infants.

Animal Data

Radiolabeled canagliflozin administered to lactating rats on day 13 post-partum was present at a milk/plasma ratio of 1.40, indicating that canagliflozin and its metabolites are transferred into milk at a concentration comparable to that in plasma. Juvenile rats directly exposed to canagliflozin showed a risk to the developing kidney (renal pelvic and tubular dilatations) during maturation.

8.3 Females and Males of Reproductive Potential

Discuss the potential for unintended pregnancy with premenopausal women as therapy with metformin may result in ovulation in some anovulatory women.

8.4 Pediatric Use

Safety and effectiveness of INVOKAMET/INVOKAMET XR in pediatric patients under 18 years of age have not been established.

8.5 Geriatric Use

INVOKAMET and INVOKAMET XR

Because renal function abnormalities can occur after initiating canagliflozin, metformin is substantially excreted by the kidney, and aging can be associated with reduced renal function, monitor renal function more frequently after initiating INVOKAMET/INVOKAMET XR in the elderly and then adjust dose based on renal function [see Dosage and Administration (2.4) and Warnings and Precautions (5.1, 5.5)].

Canagliflozin

In 13 clinical trials of canagliflozin, 2,294 patients 65 years and older, and 351 patients 75 years and older were exposed to canagliflozin. Of these patients, 1,534 patients 65 years and older and 196 patients 75 years and older were exposed to the combination of canagliflozin and metformin [see Clinical Studies (14)]. Patients 65 years and older had a higher incidence of adverse reactions related to reduced intravascular volume with canagliflozin (such as hypotension, postural dizziness, orthostatic hypotension, syncope, and dehydration), particularly with the 300 mg daily dose, compared to younger patients; a more prominent increase in the incidence was seen in patients who were 75 years and older [see Dosage and Administration (2.1) and Adverse Reactions (6.1)]. Smaller reductions in HbA_{1c} with canagliflozin relative to placebo were seen in older (65 years and older; -0.61% with canagliflozin 100 mg and -0.74% with canagliflozin 300 mg relative to placebo) compared to younger patients (-0.72% with canagliflozin 100 mg and -0.87% with canagliflozin 300 mg relative to placebo).

Metformin

Controlled clinical trials of metformin did not include sufficient numbers of elderly patients to determine whether they respond differently from younger patients, although other reported clinical experience has not identified differences in responses between the elderly and younger patients. The initial and maintenance dosing of metformin should be conservative in patients with advanced age due to the potential for decreased renal function in this population. Any dose adjustment should be based on a careful assessment of renal function [see Contraindications (4), Warnings and Precautions (5.5), and Clinical Pharmacology (12.3)].

8.6 Renal Impairment

Canagliflozin

The efficacy and safety of canagliflozin were evaluated in a trial that included patients with moderate renal impairment (eGFR 30 to less than 50 mL/min/1.73 m²). These patients had less overall glycemic efficacy and had a higher occurrence of adverse reactions related to reduced intravascular volume, renal-related adverse reactions, and decreases in eGFR compared to patients with mild renal impairment or normal renal function (eGFR greater than or equal to 60 mL/min/1.73 m²). Dose-related, transient mean increases in serum potassium

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

were observed early after initiation of canagliflozin (i.e., within 3 weeks) in this trial. Increases in serum potassium of greater than 5.4 mEq/L and 15% above baseline occurred in 16.1%, 12.4%, and 27.0% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively. Severe elevations (greater than or equal to 6.5 mEq/L) occurred in 1.1%, 2.2%, and 2.2% of patients treated with placebo, canagliflozin 100 mg, and canagliflozin 300 mg, respectively [see Dosage and Administration (2.4), Contraindications (4), Warnings and Precautions (5.3, 5.5), and Adverse Reactions (6.1)].

The efficacy and safety of canagliflozin have not been established in patients with severe renal impairment (eGFR less than 30 mL/min/1.73 m²), with ESRD, or receiving dialysis. Canagliflozin is not expected to be effective in these patient populations [see Contraindications (4) and Clinical Pharmacology (12.3)].

8.7 Hepatic Impairment

Use of metformin in patients with hepatic impairment has been associated with some cases of lactic acidosis. INVOKAMET/INVOKAMET XR is not recommended in patients with hepatic impairment [see Warnings and Precautions (5.1)].

10 OVERDOSAGE

Overdose of metformin hydrochloride has occurred, including ingestion of amounts greater than 50 grams. Hypoglycemia was reported in approximately 10% of cases, but no causal association with metformin has been established. Lactic acidosis has been reported in approximately 32% of metformin overdose cases [see Warnings and Precautions (5.1)].


In the event of an overdose with INVOKAMET/INVOKAMET XR, contact the Poison Control Center. Employ the usual supportive measures (e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring, and institute supportive treatment) as dictated by the patient's clinical status. Canagliflozin was negligibly removed during a 4-hour hemodialysis session. Canagliflozin is not expected to be dialyzable by peritoneal dialysis. Metformin is dialyzable with a clearance of up to 170 mL/min under good hemodynamic conditions. Therefore, hemodialysis may be useful partly for removal of accumulated metformin from patients in whom INVOKAMET/INVOKAMET XR overdose is suspected.

11 DESCRIPTION

INVOKAMET (canagliflozin and metformin hydrochloride) and INVOKAMET XR (canagliflozin and metformin hydrochloride extended-release) tablets contain canagliflozin and metformin hydrochloride.

Canagliflozin


Canagliflozin is an inhibitor of sodium-glucose co-transporter 2 (SGLT2), the transporter responsible for reabsorbing the majority of glucose filtered by the kidney. Canagliflozin is chemically known as (1S)-1,5-anhydro-1-[3-[[5-(4-fluorophenyl)-2-thienyl]methyl]-4-methylphenyl]-D-glucitol hemihydrate and its molecular formula and weight are C₂₄H₂₅F₀O₅S•1/2 H₂O and 453.53, respectively. The structural formula for canagliflozin is:



Canagliflozin is practically insoluble in aqueous media from pH 1.1 to 12.9.

Metformin Hydrochloride

Metformin hydrochloride is a biguanide chemically known as 1,1-Dimethylbiguanide hydrochloride and its molecular formula and weight are C₄H₁₁N₅•HCl and 165.62, respectively. The structural formula for metformin hydrochloride is:



INVOKAMET and INVOKAMET XR

INVOKAMET/INVOKAMET XR are supplied as film-coated tablets for oral administration. Each 50 mg/500 mg tablet and 50 mg/1,000 mg tablet contains 51 mg of canagliflozin equivalent to 50 mg canagliflozin (anhydrous) and 500 mg or 1,000 mg metformin hydrochloride (equivalent to metformin 389.93 mg and 779.86 mg, respectively). Each 150 mg/500 mg tablet and 150 mg/1,000 mg tablet contains 153 mg of canagliflozin equivalent to 150 mg canagliflozin (anhydrous) and 500 mg or 1,000 mg metformin hydrochloride (equivalent to metformin 389.93 mg and 779.86 mg, respectively).

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

INVOKAMET contains the following inactive ingredients: croscarmellose sodium, hypromellose, magnesium stearate, and microcrystalline cellulose. The magnesium stearate is vegetable-sourced. The tablets are finished with a commercially available film-coating consisting of the following inactive ingredients: macrogol/PEG3350, polyvinyl alcohol (partially hydrolyzed), talc, titanium dioxide, iron oxide yellow (50 mg/1,000 mg and 150 mg/500 mg tablets only), iron oxide red (50 mg/1,000 mg, 150 mg/500 mg and 150 mg/1,000 mg tablets only), and iron oxide black (150 mg/1,000 mg tablets only).

INVOKAMET XR contains the following inactive ingredients: croscarmellose sodium, hydroxypropyl cellulose, hypromellose, lactose anhydrous, magnesium stearate (vegetable-sourced), microcrystalline cellulose, polyethylene oxide, and silicified microcrystalline cellulose (50 mg/500 mg and 50 mg/1,000 mg tablets only). The tablets are finished with a commercially available film-coating consisting of the following inactive ingredients: macrogol/PEG3350, polyvinyl alcohol (partially hydrolyzed), talc, titanium dioxide, iron oxide red, iron oxide yellow, and iron oxide black (50 mg/1,000 mg and 150 mg/1,000 mg tablets only).

INVOKAMET XR tablets provide canagliflozin for immediate-release and metformin hydrochloride for extended-release. Each bilayer tablet is compressed from two separate granulates, one for each active ingredient of the tablet, and finished with a film-coating. The metformin HCl extended-release layer is based on a polymer matrix which controls the drug release by passive diffusion through the swollen matrix in combination with tablet erosion.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Canagliflozin

Sodium-glucose co-transporter 2 (SGLT2), expressed in the proximal renal tubules, is responsible for the majority of the reabsorption of filtered glucose from the tubular lumen. Canagliflozin is an inhibitor of SGLT2. By inhibiting SGLT2, canagliflozin reduces reabsorption of filtered glucose and lowers the renal threshold for glucose (RT_G), and thereby increases urinary glucose excretion (UGE).

Metformin

Metformin is an antihyperglycemic agent which improves glucose tolerance in patients with type 2 diabetes, lowering both basal and postprandial plasma glucose. Metformin decreases hepatic glucose production, decreases intestinal absorption of glucose, and improves insulin sensitivity by increasing peripheral glucose uptake and utilization. With metformin therapy, insulin secretion remains unchanged while fasting insulin levels and day-long plasma insulin response may decrease.

12.2 Pharmacodynamics

Canagliflozin

Following single and multiple oral doses of canagliflozin in patients with type 2 diabetes, dose-dependent decreases in RT_G and increases in urinary glucose excretion were observed. From a starting RT_G value of approximately 240 mg/dL, canagliflozin at 100 mg and 300 mg once daily suppressed RT_G throughout the 24-hour period. Maximal suppression of mean RT_G over the 24-hour period was seen with the 300 mg daily dose to approximately 70 to 90 mg/dL in patients with type 2 diabetes in Phase 1 trials. The reductions in RT_G led to increases in mean UGE of approximately 100 g/day in patients with type 2 diabetes treated with either 100 mg or 300 mg of canagliflozin. The 24-h mean RT_G at steady state was similar following once daily and twice daily dosing regimens at the same total daily dose of 100 mg or 300 mg. In patients with type 2 diabetes given 100 to 300 mg once daily over a 16-day dosing period, reductions in RT_G and increases in urinary glucose excretion were observed over the dosing period. In this trial, plasma glucose declined in a dose-dependent fashion within the first day of dosing.

Cardiac Electrophysiology

In a randomized, double-blind, placebo-controlled, active-comparator, 4-way crossover trial, 60 healthy subjects were administered a single oral dose of canagliflozin 300 mg, canagliflozin 1,200 mg (4 times the maximum recommended dose), moxifloxacin, and placebo. No meaningful changes in QTc interval were observed with either the recommended dose of 300 mg or the 1,200 mg dose.

12.3 Pharmacokinetics

INVOKAMET

The results of a bioequivalence trial in healthy subjects demonstrated that INVOKAMET 50 mg/500 mg, 50 mg/1,000 mg, 150 mg/500 mg and 150 mg/1,000 mg combination tablets are bioequivalent to co-administration of corresponding doses of canagliflozin and metformin HCl as individual tablets under fed conditions. Administration of INVOKAMET 150 mg/1,000 mg fixed-dose combination with food resulted in no change in overall exposure of canagliflozin. There was no change in metformin AUC; however, the mean peak plasma concentration of metformin was decreased by 16% when administered with food. A delayed time to peak plasma concentration was observed for both components (a delay of 2 hours for canagliflozin and 1 hour for metformin) under fed conditions. These changes are not likely to be clinically meaningful.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

INVOKAMET XR

After administration of INVOKAMET XR tablets with a high-fat breakfast, the peak (C_{max}) and total (AUC) exposure of canagliflozin were not altered relative to dosing in the fasted state. However, the AUC of metformin increased by approximately 61% and C_{max} increased by approximately 13%.

Canagliflozin

The pharmacokinetics of canagliflozin is essentially similar in healthy subjects and patients with type 2 diabetes. Following single-dose oral administration of 100 mg and 300 mg of canagliflozin, peak plasma concentrations (median T_{max}) of canagliflozin occurs within 1 to 2 hours post-dose. Plasma C_{max} and AUC of canagliflozin increased in a dose-proportional manner from 50 mg to 300 mg. The apparent terminal half-life (t_{1/2}) was 10.6 hours and 13.1 hours for the 100 mg and 300 mg doses, respectively. Steady-state was reached after 4 to 5 days of once-daily dosing with canagliflozin 100 mg to 300 mg. Canagliflozin does not exhibit time-dependent pharmacokinetics and accumulated in plasma up to 36% following multiple doses of 100 mg and 300 mg. The mean systemic exposure (AUC) at steady state was similar following once daily and twice daily dosing regimens at the same total daily dose of 100 mg or 300 mg.

Absorption

Canagliflozin

The mean absolute oral bioavailability of canagliflozin is approximately 65%.

Metformin

The absolute bioavailability of a metformin HCl 500 mg tablet given under fasting conditions is approximately 50% to 60%. Trials using single oral doses of metformin hydrochloride 500 to 1,500 mg, and 850 to 2,550 mg, indicate that there is a lack of dose proportionality with increasing doses, which is due to decreased absorption rather than an alteration in elimination.

Following a single oral dose of 1,000 mg (two 500 mg tablets) metformin HCl extended-release after a meal, the time to reach maximum plasma metformin concentration (T_{max}) is achieved at approximately 7-8 hours. In both single and multiple-dose trials in healthy subjects, once daily 1,000 mg (two 500 mg tablets) dosing results in up to 35% higher C_{max} of metformin relative to the immediate-release given as 500 mg twice daily without any change in overall systemic exposure, as measured by AUC.

Distribution

Canagliflozin

The mean steady-state volume of distribution of canagliflozin following a single intravenous infusion in healthy subjects was 83.5 L, suggesting extensive tissue distribution. Canagliflozin is extensively bound to proteins in plasma (99%), mainly to albumin. Protein binding is independent of canagliflozin plasma concentrations. Plasma protein binding is not meaningfully altered in patients with renal or hepatic impairment.

Metformin

The apparent volume of distribution (V/F) of metformin following single oral doses of metformin HCl 850 mg immediate-release tablets averaged 654 ± 358 L. Metformin is negligibly bound to plasma proteins, in contrast to sulfonyleureas, which are more than 90% protein bound. Metformin partitions into erythrocytes, most likely as a function of time. At usual clinical doses and dosing schedules of metformin HCl tablets, steady-state plasma concentrations of metformin are reached within 24 to 48 hours and are generally less than 1 mcg/mL.

Metabolism

Canagliflozin

O-glucuronidation is the major metabolic elimination pathway for canagliflozin, which is mainly glucuronidated by UGT1A9 and UGT2B4 to two inactive O-glucuronide metabolites. CYP3A4-mediated (oxidative) metabolism of canagliflozin is minimal (approximately 7%) in humans.

Metformin

Intravenous single-dose trials in normal subjects demonstrate that metformin is excreted unchanged in the urine and does not undergo hepatic metabolism (no metabolites have been identified in humans) or biliary excretion.

Excretion

Canagliflozin

Following administration of a single oral [¹⁴C] canagliflozin dose to healthy subjects, 41.5%, 7.0%, and 3.2% of the administered radioactive dose was recovered in feces as canagliflozin, a hydroxylated metabolite, and an O-glucuronide metabolite, respectively. Enterohepatic circulation of canagliflozin was negligible.

Approximately 33% of the administered radioactive dose was excreted in urine, mainly as O-glucuronide metabolites (30.5%). Less than 1% of the dose was excreted as unchanged canagliflozin in urine. Renal clearance of canagliflozin 100 mg and 300 mg doses ranged from 1.30 to 1.55 mL/min.

Mean systemic clearance of canagliflozin was approximately 192 mL/min in healthy subjects following intravenous administration.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Metformin

Renal clearance is approximately 3.5 times greater than creatinine clearance, which indicates that tubular secretion is the major route of metformin elimination. Following oral administration, approximately 90% of the absorbed drug is eliminated via the renal route within the first 24 hours, with a plasma elimination half-life of approximately 6.2 hours. In blood, the elimination half-life is approximately 17.6 hours, suggesting that the erythrocyte mass may be a compartment of distribution.

Specific Populations

Trials characterizing the pharmacokinetics of canagliflozin and metformin after administration of INVOKAMET/INVOKAMET XR were not conducted in patients with renal and hepatic impairment. Descriptions of the individual components in this patient population are described below.

Renal Impairment

Canagliflozin

A single-dose, open-label trial evaluated the pharmacokinetics of canagliflozin 200 mg in subjects with varying degrees of renal impairment (classified using the MDRD-eGFR formula) compared to healthy subjects.

Renal impairment did not affect the C_{max} of canagliflozin. Compared to healthy subjects (N=3; eGFR greater than or equal to 90 mL/min/1.73 m²), plasma AUC of canagliflozin was increased by approximately 15%, 29%, and 53% in subjects with mild (N=10), moderate (N=9), and severe (N=10) renal impairment, respectively, (eGFR 60 to less than 90, 30 to less than 60, and 15 to less than 30 mL/min/1.73 m², respectively), but was similar for ESRD (N=8) subjects and healthy subjects. Increases in canagliflozin AUC of this magnitude are not considered clinically relevant. The pharmacodynamic response to canagliflozin declines with increasing severity of renal impairment [see *Contraindications (4) and Warnings and Precautions (5.5)*].

Canagliflozin was negligibly removed by hemodialysis.

Metformin

In patients with decreased renal function, the plasma and blood half-life of metformin is prolonged and the renal clearance is decreased [see *Contraindications (4) and Warnings and Precautions (5.1)*].

Following a single dose administration of metformin HCl extended-release 500 mg in patients with mild and moderate renal failure (based on measured creatinine clearance), the oral and renal clearance of metformin were decreased by 33% and 50% and 16% and 53%, respectively [see *Warnings and Precautions (5.5)*]. Metformin peak and systemic exposure was 27% and 61% greater, respectively in mild renal impaired and 74% and 2.36-fold greater in moderate renal impaired patients as compared to healthy subjects [see *Contraindications (4) and Warnings and Precautions (5.1)*].

Hepatic Impairment

Canagliflozin

Relative to subjects with normal hepatic function, the geometric mean ratios for C_{max} and AUC_∞ of canagliflozin were 107% and 110%, respectively, in subjects with Child-Pugh class A (mild hepatic impairment) and 96% and 111%, respectively, in subjects with Child-Pugh class B (moderate hepatic impairment) following administration of a single 300 mg dose of canagliflozin.

These differences are not considered to be clinically meaningful. There is no clinical experience in patients with Child-Pugh class C (severe) hepatic impairment [see *Warnings and Precautions (5.1)*].

Metformin

No pharmacokinetic trials of metformin have been conducted in patients with hepatic insufficiency [see *Warnings and Precautions (5.1)*].

Pharmacokinetic Effects of Age, Body Mass Index (BMI)/Weight, Gender and Race
Canagliflozin

Based on the population PK analysis with data collected from 1526 subjects, age, body mass index (BMI)/weight, gender, and race do not have a clinically meaningful effect on the pharmacokinetics of canagliflozin [see *Use in Specific Populations (8.5)*].

Metformin

Metformin pharmacokinetic parameters did not differ significantly between normal subjects and patients with type 2 diabetes when analyzed according to gender.

No trials of metformin pharmacokinetic parameters according to race have been performed.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Canagliflozin

Age had no clinically meaningful effect on the pharmacokinetics of canagliflozin based on a population pharmacokinetic analysis [see *Adverse Reactions (6.1) and Use in Specific Populations (8.5)*].

Metformin

Limited data from controlled pharmacokinetic trials of metformin in healthy elderly subjects suggest that total plasma clearance of metformin is decreased, the half-life is prolonged, and C_{max} is increased, compared with healthy young subjects. From these data, it appears that the change in metformin pharmacokinetics with aging is primarily accounted for by a change in renal function [see *Warnings and Precautions (5.1, 5.5) and Use in Specific Populations (8.5)*].

Drug-Drug Interactions

INVOKAMET and INVOKAMET XR

Pharmacokinetic drug interaction trials with INVOKAMET/INVOKAMET XR have not been performed; however, such trials have been conducted with the individual components canagliflozin and metformin hydrochloride.

Co-administration of multiple doses of canagliflozin (300 mg) and metformin (2,000 mg) given once daily did not meaningfully alter the pharmacokinetics of either canagliflozin or metformin in healthy subjects.

Canagliflozin

In Vitro Assessment of Drug Interactions

Canagliflozin did not induce CYP450 enzyme expression (3A4, 2C9, 2C19, 2B6, and 1A2) in cultured human hepatocytes. Canagliflozin did not inhibit the CYP450 isoenzymes (1A2, 2A6, 2C19, 2D6, or 2E1) and weakly inhibited CYP2B6, CYP2C8, CYP2C9, and CYP3A4 based on *in vitro* studies with human hepatic microsomes. Canagliflozin is a weak inhibitor of P-gp.

Canagliflozin is also a substrate of drug transporters P-glycoprotein (P-gp) and MRP2.

In Vivo Assessment of Drug Interactions

Table 7: Effect of Co-Administered Drugs on Systemic Exposures of Canagliflozin

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Canagliflozin*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.0	
			AUC [†] (90% CI)	C _{max} (90% CI)
See Drug Interactions (7.2) for the clinical relevance of the following:				
Rifampin	600 mg QD for 8 days	300 mg	0.49 (0.44; 0.54)	0.72 (0.61; 0.84)
No dose adjustments of canagliflozin required for the following:				
Cyclosporine	400 mg	300 mg QD for 8 days	1.23 (1.19; 1.27)	1.01 (0.91; 1.11)
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg QD for 6 days	0.91 (0.88; 0.94)	0.92 (0.84; 0.99)
Hydrochlorothiazide	25 mg QD for 35 days	300 mg QD for 7 days	1.12 (1.08; 1.17)	1.15 (1.06; 1.25)
Metformin	2,000 mg	300 mg QD for 8 days	1.10 (1.05; 1.15)	1.05 (0.96; 1.16)
Probenecid	500 mg BID for 3 days	300 mg QD for 17 days	1.21 (1.16; 1.25)	1.13 (1.00; 1.28)

* Single dose unless otherwise noted

† AUC_{inf} for drugs given as a single dose and AUC_{24h} for drugs given as multiple doses

QD = once daily; BID = twice daily

Table 8: Effect of Canagliflozin on Systemic Exposure of Co-Administered Drugs

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Canagliflozin*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.0		
			AUC [†] (90% CI)	C _{max} (90% CI)	
See Drug Interactions (7.2) for the clinical relevance of the following:					
Digoxin	0.5 mg QD first day followed by 0.25 mg QD for 6 days	300 mg QD for 7 days	Digoxin	1.20 (1.12; 1.28)	1.36 (1.21; 1.53)
No dose adjustments of co-administered drug required for the following:					
Acetaminophen	1,000 mg	300 mg BID for 25 days	Acetaminophen	1.06 [‡] (0.98; 1.14)	1.00 (0.92; 1.09)
Ethinyl estradiol and levonorgestrel	0.03 mg ethinyl estradiol and 0.15 mg levonorgestrel	200 mg QD for 6 days	ethinyl estradiol	1.07 (0.99; 1.15)	1.22 (1.10; 1.35)
			Levonorgestrel	1.06 (1.00; 1.13)	1.22 (1.11; 1.35)
Glyburide	1.25 mg	200 mg QD for 6 days	Glyburide	1.02 (0.98; 1.07)	0.93 (0.85; 1.01)
			3-cis-hydroxy-glyburide	1.01 (0.96; 1.07)	0.99 (0.91; 1.08)
			4-trans-hydroxy-glyburide	1.03 (0.97; 1.09)	0.96 (0.88; 1.04)
Hydrochlorothiazide	25 mg QD for 35 days	300 mg QD for 7 days	Hydrochlorothiazide	0.99 (0.95; 1.04)	0.94 (0.87; 1.01)
Metformin	2,000 mg	300 mg QD for 8 days	Metformin	1.20 (1.08; 1.34)	1.06 (0.93; 1.20)
Simvastatin	40 mg	300 mg QD for 7 days	Simvastatin	1.12 (0.94; 1.33)	1.09 (0.91; 1.31)
			simvastatin acid	1.18 (1.03; 1.35)	1.26 (1.10; 1.45)
Warfarin	30 mg	300 mg QD for 12 days	(R)-warfarin	1.01 (0.96; 1.06)	1.03 (0.94; 1.13)
			(S)-warfarin	1.06 (1.00; 1.12)	1.01 (0.90; 1.13)
			INR	1.00 (0.98; 1.03)	1.05 (0.99; 1.12)

* Single dose unless otherwise noted
[†] AUC_{inf} for drugs given as a single dose and AUC_{24h} for drugs given as multiple doses
[‡] AUC_{0-12h}
 QD = once daily; BID = twice daily; INR = International Normalized Ratio

Metformin

Table 9: Effect of Co-Administered Drugs on Plasma Metformin Systemic Exposures

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Metformin HCl*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.00	
			AUC [†]	C _{max}
No dose adjustments required for the following:				
Glyburide	5 mg	500 mg [‡]	0.98 [§]	0.99 [§]
Furosemide	40 mg	850 mg	1.09 [§]	1.22 [§]
Nifedipine	10 mg	850 mg	1.16	1.21
Propranolol	40 mg	850 mg	0.90	0.94
Ibuprofen	400 mg	850 mg	1.05 [§]	1.07 [§]
Drugs that are eliminated by renal tubular secretion increase the accumulation of metformin [see Warnings and Precautions (5) and Drug Interactions (7)]				
Cimetidine	400 mg	850 mg	1.40	1.61
Carbonic anhydrase inhibitors may cause metabolic acidosis [see Warnings and Precautions (5) and Drug Interactions (7)]				
Topiramate [¶]	100 mg	500 mg	1.25 [#]	1.18

* Single dose unless otherwise noted
[†] AUC = AUC_{0-∞}
[‡] Metformin HCl extended-release tablets 500 mg
[§] Ratio of arithmetic means
[¶] Healthy volunteer study at steady state with topiramate 100 mg every 12 hours and metformin 500 mg every 12 hours for 7 days. Study conducted to assess pharmacokinetics only
[#] Steady state AUC_{0-12h}.

Table 10: Effect of Metformin on Co-Administered Drug Systemic Exposures

Co-Administered Drug	Dose of Co-Administered Drug*	Dose of Metformin HCl*	Geometric Mean Ratio (Ratio With/Without Co-Administered Drug) No Effect = 1.00	
			AUC [†]	C _{max}
No dose adjustments required for the following:				
Glyburide	5 mg	500 mg [‡]	0.78 [§]	0.63 [§]
Furosemide	40 mg	850 mg	0.87 [§]	0.69 [§]
Nifedipine	10 mg	850 mg	1.10 [‡]	1.08
Propranolol	40 mg	850 mg	1.01 [‡]	0.94
Ibuprofen	400 mg	850 mg	0.97 [¶]	1.01 [¶]
Cimetidine	400 mg	850 mg	0.95 [‡]	1.01

* Single dose unless otherwise noted
[†] AUC = AUC_{0-∞}
[‡] AUC_{0-24 hr} reported
[§] Ratio of arithmetic means, p-value of difference <0.05
[¶] Ratio of arithmetic means.

Metformin is negligibly bound to plasma proteins and is, therefore, less likely to interact with highly protein-bound drugs such as salicylates, sulfonamides, chloramphenicol, and probenecid.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

INVOKAMET and INVOKAMET XR

No animal studies have been conducted with the combined products in INVOKAMET/INVOKAMET XR to evaluate carcinogenesis, mutagenesis, or impairment of fertility. The following data are based on findings in studies with canagliflozin and metformin individually.

Canagliflozin

Carcinogenesis

Carcinogenicity was evaluated in 2-year studies conducted in CD1 mice and Sprague-Dawley rats. Canagliflozin did not increase the incidence of tumors in mice dosed at 10, 30, or 100 mg/kg (less than or equal to 14 times exposure from a 300 mg clinical dose).

Testicular Leydig cell tumors, considered secondary to increased luteinizing hormone (LH), increased significantly in male rats at all doses tested (10, 30, and 100 mg/kg). In a 12-week clinical trial, LH did not increase in males treated with canagliflozin.

Renal tubular adenoma and carcinoma increased significantly in male and female rats dosed at 100 mg/kg, or approximately 12-times exposure from a 300 mg clinical dose. Also, adrenal pheochromocytoma increased significantly in males and numerically in females dosed at 100 mg/kg. Carbohydrate malabsorption associated with high doses of canagliflozin was considered a necessary proximal event in the emergence of renal and adrenal tumors in rats. Clinical trials have not demonstrated carbohydrate malabsorption in humans at canagliflozin doses of up to 2-times the recommended clinical dose of 300 mg.

Mutagenesis

Canagliflozin was not mutagenic with or without metabolic activation in the Ames assay. Canagliflozin was mutagenic in the *in vitro* mouse lymphoma assay with but not without metabolic activation. Canagliflozin was not mutagenic or clastogenic in an *in vivo* oral micronucleus assay in rats and an *in vivo* oral Comet assay in rats.

Metformin

Carcinogenesis

Long-term carcinogenicity studies have been performed in rats (dosing duration of 104 weeks) and mice (dosing duration of 91 weeks) at doses up to and including 900 mg/kg/day and 1500 mg/kg/day, respectively. These doses are both approximately 4 times the maximum recommended human daily dose of 2000 mg based on body surface area comparisons. No evidence of carcinogenicity with metformin was found in either male or female mice. Similarly, there was no tumorigenic potential observed with metformin in male rats. There was, however, an increased incidence of benign stromal uterine polyps in female rats treated with 900 mg/kg/day.

Mutagenesis

There was no evidence of a mutagenic potential of metformin in the following *in vitro* tests: Ames test (*S. typhimurium*), gene mutation test (mouse lymphoma cells), or chromosomal aberrations test (human lymphocytes). Results in the *in vivo* mouse micronucleus test were also negative.

Impairment of Fertility

Canagliflozin had no effects on the ability of rats to mate and sire or maintain a litter up to the high dose of 100 mg/kg (approximately 14 times and 18 times the 300 mg clinical dose in males and females, respectively), although there were minor alterations in a number of reproductive parameters (decreased sperm velocity, increased number of abnormal sperm, slightly fewer corpora lutea, fewer implantation sites, and smaller litter sizes) at the highest dosage administered.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Fertility of male or female rats was unaffected by metformin when administered at doses as high as 600 mg/kg/day, which is approximately 3 times the maximum recommended human daily dose based on body surface area comparisons.

14 CLINICAL STUDIES

Canagliflozin has been studied in combination with metformin alone, metformin and sulfonyleurea, metformin and sitagliptin, metformin and a thiazolidinedione (i.e., pioglitazone), and metformin and insulin (with or without other anti-hyperglycemic agents). The efficacy of canagliflozin was compared to a dipeptidyl peptidase-4 (DPP-4) inhibitor (sitagliptin), both as add-on combination therapy with metformin and sulfonyleurea, and a sulfonyleurea (glimepiride), both as add-on combination therapy with metformin.

There have been no clinical efficacy trials conducted with INVOKAMET/INVOKAMET XR; however, bioequivalence of INVOKAMET/INVOKAMET XR to canagliflozin and metformin HCl co-administered as individual tablets was demonstrated in healthy subjects.

In patients with type 2 diabetes, treatment with canagliflozin and metformin produced clinically and statistically significant improvements in HbA_{1c} compared to placebo. Reductions in HbA_{1c} were observed across subgroups including age, gender, race, and baseline body mass index (BMI).

14.1 Glycemic Control Trials in Adults with Type 2 Diabetes Mellitus

Canagliflozin as Initial Combination Therapy with Metformin

A total of 1,186 patients with type 2 diabetes inadequately controlled with diet and exercise participated in a 26-week double-blind, active-controlled, parallel-group, 5-arm, multicenter trial to evaluate the efficacy and safety of initial therapy with canagliflozin in combination with metformin XR. The median age was 56 years, 48% of patients were men, and the mean baseline eGFR was 87.6 mL/min/1.73 m². The median duration of diabetes was 1.6 years, and 72% of patients were treatment naïve. After completing a 2-week single-blind placebo run-in period, patients were randomly assigned for a double-blind treatment period of 26 weeks to 1 of 5 treatment groups (Table 11). The metformin HCl XR dose was initiated at 500 mg/day for the first week of treatment and then increased to 1,000 mg/day. Metformin HCl XR or matching placebo was up-titrated every 2-3 weeks during the next 8 weeks of treatment to a maximum daily dose of 1,500 to 2,000 mg/day, as tolerated; about 90% of patients reached 2,000 mg/day.

At the end of treatment, canagliflozin 100 mg and canagliflozin 300 mg in combination with metformin HCl XR resulted in a statistically significant greater improvement in HbA_{1c} compared to their respective canagliflozin doses (100 mg and 300 mg) alone or metformin HCl XR alone.

Table 11: Results from 26-Week Active-Controlled Clinical Study of Canagliflozin Alone or Canagliflozin as Initial Combination Therapy with Metformin HCl*

Efficacy Parameter	Metformin XR (N=237)	Canagliflozin 100 mg (N=237)	Canagliflozin 300 mg (N=238)	Canagliflozin 100 mg + Metformin HCl XR (N=237)	Canagliflozin 300 mg + Metformin HCl XR (N=237)
HbA_{1c} (%)					
Baseline (mean)	8.81	8.78	8.77	8.83	8.90
Change from baseline (adjusted mean) [†]	-1.30	-1.37	-1.42	-1.77	-1.78
Difference from canagliflozin 100 mg (adjusted mean) (95% CI) [‡]				-0.40 [‡] (-0.59, -0.21)	
Difference from canagliflozin 300 mg (adjusted mean) (95% CI) [‡]					-0.36 [‡] (-0.56, -0.17)
Difference from metformin XR (adjusted mean) (95% CI) [‡]				-0.46 [‡] (-0.66, -0.27)	-0.48 [‡] (-0.67, -0.28)
Percent of patients achieving HbA_{1c} < 7%	38	34	39	47 ^{§§}	51 ^{§§}

* Intent-to-treat population

[†] Least squares mean adjusted for covariates including baseline value and stratification factor

[‡] Adjusted p=0.001

^{§§} Adjusted p<0.05

[¶] There were 121 patients without week 26 efficacy data. Analyses addressing missing data gave consistent results with the results provided in this table.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Canagliflozin as Add-on Combination Therapy with Metformin

A total of 1,284 patients with type 2 diabetes inadequately controlled on metformin HCl monotherapy (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) participated in a 26-week, double-blind, placebo- and active-controlled trial to evaluate the efficacy and safety of canagliflozin in combination with metformin. The mean age was 55 years, 47% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m². Patients already on the required metformin HCl dose (N=1009) were randomized after completing a 2-week, single-blind, placebo run-in period. Patients taking less than the required metformin HCl dose or patients on metformin HCl in combination with another antihyperglycemic agent (N=275) were switched to metformin HCl monotherapy (at doses described above) for at least 8 weeks before entering the 2-week, single-blind, placebo run-in. After the placebo run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, sitagliptin 100 mg, or placebo, administered once daily as add-on therapy to metformin HCl.

At the end of treatment, canagliflozin 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1c} (p<0.001 for both doses) compared to placebo when added to metformin HCl. Canagliflozin 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1c} less than 7%, in significant reduction in fasting plasma glucose (FPG), in improved postprandial glucose (PPG), and in percent body weight reduction compared to placebo when added to metformin HCl (see Table 12). Statistically significant (p<0.001 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -5.4 mmHg and -6.6 mmHg with canagliflozin 100 mg and 300 mg, respectively.

Table 12: Results from 26-Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin HCl*

Efficacy Parameter	Placebo + Metformin HCl (N=183)	Canagliflozin 100 mg + Metformin HCl (N=368)	Canagliflozin 300 mg + Metformin HCl (N=367)
HbA_{1c} (%)			
Baseline (mean)	7.96	7.94	7.95
Change from baseline (adjusted mean)	-0.17	-0.79	-0.94
Difference from placebo (adjusted mean) (95% CI) [†]		-0.62 [‡] (-0.76, -0.48)	-0.77 [‡] (-0.91, -0.64)
Percent of patients achieving HbA_{1c} < 7%	30	46 [‡]	58 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	164	169	173
Change from baseline (adjusted mean)	2	-27	-38
Difference from placebo (adjusted mean) (95% CI) [†]		-30 [‡] (-36, -24)	-40 [‡] (-46, -34)
2-hour Postprandial Glucose (mg/dL)			
Baseline (mean)	249	258	262
Change from baseline (adjusted mean)	-10	-48	-57
Difference from placebo (adjusted mean) (95% CI) [†]		-38 [‡] (-49, -27)	-47 [‡] (-58, -36)
Body Weight			
Baseline (mean) in kg	86.7	88.7	85.4
% change from baseline (adjusted mean)	-1.2	-3.7	-4.2
Difference from placebo (adjusted mean) (95% CI) [†]		-2.5 [‡] (-3.1, -1.9)	-2.9 [‡] (-3.5, -2.3)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] p<0.001

Canagliflozin Compared to Glimepiride, Both as Add-on Combination Therapy with Metformin

A total of 1,450 patients with type 2 diabetes inadequately controlled on metformin HCl monotherapy (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) participated in a 52-week, double-blind, active-controlled trial to evaluate the efficacy and safety of canagliflozin in combination with metformin.

The mean age was 56 years, 52% of patients were men, and the mean baseline eGFR was 90 mL/min/1.73 m². Patients tolerating maximally required metformin HCl dose (N=928) were randomized after completing a 2-week, single-blind, placebo run-in period. Other patients (N=522) were switched to metformin HCl monotherapy (at doses described above) for at least 10 weeks, then completed a

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

2-week single-blind run-in period. After the 2-week run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, or glimepiride (titration allowed throughout the 52-week trial to 6 or 8 mg), administered once daily as add-on therapy to metformin HCl.

As shown in Table 13 and Figure 1, at the end of treatment, canagliflozin 100 mg provided similar reductions in HbA_{1c} from baseline compared to glimepiride when added to metformin HCl therapy. Canagliflozin 300 mg provided a greater reduction from baseline in HbA_{1c} compared to glimepiride, and the relative treatment difference was -0.12% (95% CI: -0.22; -0.02). As shown in Table 13, treatment with canagliflozin 100 mg and 300 mg daily provided greater improvements in percent body weight change, relative to glimepiride.

Table 13: Results from 52-Week Clinical Study Comparing Canagliflozin to Glimepiride in Combination with Metformin*

Efficacy Parameter	Canagliflozin 100 mg + Metformin HCl (N=483)	Canagliflozin 300 mg + Metformin HCl (N=485)	Glimepiride (titrated) + Metformin HCl (N=482)
HbA_{1c} (%)			
Baseline (mean)	7.78	7.79	7.83
Change from baseline (adjusted mean)	-0.82	-0.93	-0.81
Difference from glimepiride (adjusted mean) (95% CI) [†]	-0.01 [‡] (-0.11, 0.09)	-0.12 [‡] (-0.22, -0.02)	
Percent of patients achieving HbA_{1c} < 7%	54	60	56
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	165	164	166
Change from baseline (adjusted mean)	-24	-28	-18
Difference from glimepiride (adjusted mean) (95% CI) [†]	-6 (-10, -2)	-9 (-13, -5)	
Body Weight			
Baseline (mean) in kg	86.8	86.6	86.6
% change from baseline (adjusted mean)	-4.2	-4.7	1.0
Difference from glimepiride (adjusted mean) (95% CI) [†]	-5.2 [§] (-5.7, -4.7)	-5.7 [§] (-6.2, -5.1)	


* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] Canagliflozin + metformin HCl is considered non-inferior to glimepiride + metformin because the upper limit of this confidence interval is less than the pre-specified non-inferiority margin of < 0.3%.

[§] p<0.001

Figure 1: Mean HbA_{1c} Change at Each Time Point (Completers) and at Week 52 Using Last Observation Carried Forward (mITT Population)



Canagliflozin as Add-on Combination Therapy with Metformin and Sitagliptin

A total of 217 patients with type 2 diabetes inadequately controlled on the combination of metformin HCl (greater than or equal to 1,500 mg/day) and sitagliptin 100 mg/day (or equivalent fixed-dose combination) participated in a 26-week, double-blind, placebo-controlled trial to evaluate the efficacy and safety of canagliflozin in combination with metformin and sitagliptin. The mean

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

age was 57 years, 58% of patients were men, 73% of patients were Caucasian, 15% were Asian, and 12% were Black or African-American. The mean baseline eGFR was 90 mL/min/1.73 m² and the mean baseline BMI was 32 kg/m². The mean duration of diabetes was 10 years. Eligible patients entered a 2-week, single-blind, placebo run-in period and were subsequently randomized to canagliflozin 100 mg or placebo, administered once daily as add-on to metformin HCl and sitagliptin. Patients with a baseline eGFR of 70 mL/min/1.73 m² or greater who were tolerating canagliflozin 100 mg and who required additional glycemic control (fasting finger stick 100 mg/dL or greater at least twice within 2 weeks) were up-titrated to canagliflozin 300 mg. While up-titration occurred as early as Week 4, most (90%) patients randomized to canagliflozin were up-titrated to canagliflozin 300 mg by 6 to 8 weeks.

At the end of 26 weeks, canagliflozin once daily resulted in a statistically significant improvement in HbA_{1c} (p<0.001) compared to placebo when added to metformin HCl and sitagliptin (see Table 14).

Table 14: Results from 26-Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin HCl and Sitagliptin

Efficacy Parameter	Placebo + Metformin HCl and Sitagliptin (N=108*)	Canagliflozin + Metformin HCl and Sitagliptin (N=109*)
HbA_{1c} (%)		
Baseline (mean)	8.40	8.50
Change from baseline (adjusted mean)	-0.03	-0.83
Difference from placebo (adjusted mean) (95% CI) ^{†§}		-0.81 [#] (-1.11; -0.51)
Percent of patients achieving HbA_{1c} < 7%[‡]	9	28
Fasting Plasma Glucose (mg/dL)[¶]		
Baseline (mean)	180	185
Change from baseline (adjusted mean)	-3	-28
Difference from placebo (adjusted mean) (95% CI)		-25 [#] (-39; -11)

* To preserve the integrity of randomization, all randomized patients were included in the analysis. The patient who was randomized once to each arm was analyzed on canagliflozin.

[†] Early treatment discontinuation before week 26, occurred in 11.0% and 24.1% of canagliflozin and placebo patients, respectively.

[‡] Patients without week 26 efficacy data were considered as non-responders when estimating the proportion achieving HbA_{1c} < 7%.

[§] Estimated using a multiple imputation method modeling a “wash-out” of the treatment effect for patients having missing data who discontinued treatment. Missing data was imputed only at week 26 and analyzed using ANCOVA.

[¶] Estimated using a multiple imputation method modeling a “wash-out” of the treatment effect for patients having missing data who discontinued treatment. A mixed model for repeated measures was used to analyze the imputed data.

[#] p<0.001

Canagliflozin as Add-on Combination Therapy with Metformin and Sulfonylurea

A total of 469 patients with type 2 diabetes inadequately controlled on the combination of metformin HCl (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonylurea (maximal or near-maximal effective dose) participated in a 26-week, double-blind, placebo-controlled trial to evaluate the efficacy and safety of canagliflozin in combination with metformin and sulfonylurea. The mean age was 57 years, 51% of patients were men, and the mean baseline eGFR was 89 mL/min/1.73 m². Patients already on the protocol-specified doses of metformin HCl and sulfonylurea (N=372) entered a 2-week, single-blind, placebo run-in period. Other patients (N=97) were required to be on a stable protocol-specified dose of metformin HCl and sulfonylurea for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, or placebo administered once daily as add-on to metformin HCl and sulfonylurea.

At the end of treatment, canagliflozin 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1c} (p<0.001 for both doses) compared to placebo when added to metformin HCl and sulfonylurea. Canagliflozin 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1c} less than 7.0%, in a significant reduction in fasting plasma glucose (FPG), and in percent body weight reduction compared to placebo when added to metformin HCl and sulfonylurea (see Table 15).

Table 15: Results from 26-Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin and Sulfonylurea*

Efficacy Parameter	Placebo + Metformin HCl and Sulfonylurea (N=156)	Canagliflozin 100 mg + Metformin HCl and Sulfonylurea (N=157)	Canagliflozin 300 mg + Metformin HCl and Sulfonylurea (N=156)
HbA_{1c} (%)			
Baseline (mean)	8.12	8.13	8.13
Change from baseline (adjusted mean)	-0.13	-0.85	-1.06
Difference from placebo (adjusted mean) (95% CI) [†]		-0.71 [‡] (-0.90, -0.52)	-0.92 [‡] (-1.11, -0.73)
Percent of patients achieving HbA_{1c} < 7%	18	43 [‡]	57 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	170	173	168
Change from baseline (adjusted mean)	4	-18	-31
Difference from placebo (adjusted mean) (95% CI) [†]		-22 [‡] (-31, -13)	-35 [‡] (-44, -25)
Body Weight			
Baseline (mean) in kg	90.8	93.5	93.5
% change from baseline (adjusted mean)	-0.7	-2.1	-2.6
Difference from placebo (adjusted mean) (95% CI) [†]		-1.4 [‡] (-2.1, -0.7)	-2.0 [‡] (-2.7, -1.3)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy
[†] Least squares mean adjusted for baseline value and stratification factors
[‡] p<0.001

Canagliflozin Compared to Sitagliptin, Both as Add-on Combination Therapy with Metformin and Sulfonylurea

A total of 755 patients with type 2 diabetes inadequately controlled on the combination of metformin HCl (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and sulfonylurea (near-maximal or maximal effective dose) participated in a 52 week, double-blind, active-controlled trial to compare the efficacy and safety of canagliflozin 300 mg versus sitagliptin 100 mg in combination with metformin HCl and sulfonylurea. The mean age was 57 years, 56% of patients were men, and the mean baseline eGFR was 88 mL/min/1.73 m². Patients already on protocol-specified doses of metformin HCl and sulfonylurea (N=716) entered a 2-week single-blind, placebo run-in period. Other patients (N=39) were required to be on a stable protocol-specified dose of metformin HCl and sulfonylurea for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to canagliflozin 300 mg or sitagliptin 100 mg as add-on to metformin HCl and sulfonylurea.


As shown in Table 16 and Figure 2, at the end of treatment, canagliflozin 300 mg provided greater HbA_{1c} reduction compared to sitagliptin 100 mg when added to metformin HCl and sulfonylurea (p<0.05). Canagliflozin 300 mg resulted in a mean percent change in body weight from baseline of -2.5% compared to +0.3% with sitagliptin 100 mg. A mean change in systolic blood pressure from baseline of -5.06 mmHg was observed with canagliflozin 300 mg compared to +0.85 mmHg with sitagliptin 100 mg.

Table 16: Results from 52-Week Clinical Study Comparing Canagliflozin to Sitagliptin in Combination with Metformin HCl and Sulfonylurea*

Efficacy Parameter	Canagliflozin 300 mg + Metformin HCl and Sulfonylurea (N=377)	Sitagliptin 100 mg + Metformin HCl and Sulfonylurea (N=378)
HbA_{1c} (%)		
Baseline (mean)	8.12	8.13
Change from baseline (adjusted mean)	-1.03	-0.66
Difference from sitagliptin (adjusted mean) (95% CI) [†]	-0.37 [‡] (-0.50, -0.25)	
Percent of patients achieving HbA_{1c} < 7%	48	35
Fasting Plasma Glucose (mg/dL)		
Baseline (mean)	170	164
Change from baseline (adjusted mean)	-30	-6
Difference from sitagliptin (adjusted mean) (95% CI) [†]	-24 (-30, -18)	
Body Weight		
Baseline (mean) in kg	87.6	89.6
% change from baseline (adjusted mean)	-2.5	0.3
Difference from sitagliptin (adjusted mean) (95% CI) [†]	-2.8 [§] (-3.3, -2.2)	

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy
[†] Least squares mean adjusted for baseline value and stratification factors
[‡] Canagliflozin + metformin HCl + sulfonylurea is considered non-inferior to sitagliptin + metformin HCl + sulfonylurea because the upper limit of this confidence interval is less than the pre-specified non-inferiority margin of < 0.3%.
[§] p<0.001

Figure 2: Mean HbA_{1c} Change at Each Time Point (Completers) and at Week 52 Using Last Observation Carried Forward (mITT Population)



Canagliflozin as Add-on Combination Therapy with Metformin and Pioglitazone
 A total of 342 patients with type 2 diabetes inadequately controlled on the combination of metformin (greater than or equal to 2,000 mg/day or at least 1,500 mg/day if higher dose not tolerated) and pioglitazone (30 or 45 mg/day) participated in a 26-week, double-blind, placebo-controlled trial to evaluate the efficacy and safety of canagliflozin in combination with metformin and pioglitazone. The mean age was 57 years, 63% of patients were men, and the mean baseline eGFR was 86 mL/min/1.73 m². Patients already on protocol-specified doses of metformin HCl and pioglitazone (N=163) entered a 2-week, single-blind, placebo run-in period. Other patients (N=181) were required to be on stable protocol-specified doses of metformin HCl and pioglitazone for at least 8 weeks before entering the 2-week run-in period. Following the run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, or placebo, administered once daily as add-on to metformin HCl and pioglitazone.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

At the end of treatment, canagliflozin 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1c} (p<0.001 for both doses) compared to placebo when added to metformin HCl and pioglitazone. Canagliflozin 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1c} less than 7%, in significant reduction in fasting plasma glucose (FPG), and in percent body weight reduction compared to placebo when added to metformin HCl and pioglitazone (see Table 17). Statistically significant (p<0.05 for both doses) mean changes from baseline in systolic blood pressure relative to placebo were -4.1 mmHg and -3.5 mmHg with canagliflozin 100 mg and 300 mg, respectively.

Table 17: Results from 26–Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin HCl and Pioglitazone*

Efficacy Parameter	Placebo + Metformin HCl and Pioglitazone (N=115)	Canagliflozin 100 mg + Metformin HCl and Pioglitazone (N=113)	Canagliflozin 300 mg + Metformin HCl and Pioglitazone (N=114)
HbA_{1c} (%)			
Baseline (mean)	8.00	7.99	7.84
Change from baseline (adjusted mean)	-0.26	-0.89	-1.03
Difference from placebo (adjusted mean) (95% CI) [†]		-0.62 [‡] (-0.81, -0.44)	-0.76 [‡] (-0.95, -0.58)
Percent of patients achieving HbA _{1c} < 7%	33	47 [‡]	64 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline (mean)	164	169	164
Change from baseline (adjusted mean)	3	-27	-33
Difference from placebo (adjusted mean) (95% CI) [†]		-29 [‡] (-37, -22)	-36 [‡] (-43, -28)
Body Weight			
Baseline (mean) in kg	94.0	94.2	94.4
% change from baseline (adjusted mean)	-0.1	-2.8	-3.8
Difference from placebo (adjusted mean) (95% CI) [†]		-2.7 [‡] (-3.6, -1.8)	-3.7 [‡] (-4.6, -2.8)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] p<0.001

Canagliflozin as Add-on Combination Therapy with Insulin (With or Without Other Anti-Hyperglycemic Agents, Including Metformin)

A total of 1,718 patients with type 2 diabetes inadequately controlled on insulin greater than or equal to 30 units/day or insulin in combination with other antihyperglycemic agents participated in an 18-week, double-blind, placebo-controlled substudy of a cardiovascular trial to evaluate the efficacy and safety of canagliflozin in combination with insulin. Of these patients, a subgroup of 432 patients with inadequate glycemic control received canagliflozin or placebo plus metformin HCl and ≥ 30 units/day of insulin over 18 weeks.

In this subgroup, the mean age was 61 years, 67% of patients were men, and the mean baseline eGFR was 81 mL/min/1.73 m². Patients on metformin HCl in combination with basal, bolus, or basal/bolus insulin for at least 10 weeks entered a 2-week, single-blind, placebo run-in period. Approximately 74% of these patients were on a background of metformin HCl and basal/bolus insulin regimen. After the run-in period, patients were randomized to canagliflozin 100 mg, canagliflozin 300 mg, or placebo, administered once daily as add-on to metformin HCl and insulin. The mean daily insulin dose at baseline was 93 units, which was similar across treatment groups.

At the end of treatment, canagliflozin 100 mg and 300 mg once daily resulted in a statistically significant improvement in HbA_{1c} (p<0.001 for both doses) compared to placebo when added to metformin HCl and insulin. Canagliflozin 100 mg and 300 mg once daily also resulted in a greater proportion of patients achieving an HbA_{1c} less than 7%, in significant reductions in fasting plasma glucose (FPG), and in percent body weight reductions compared to placebo (see Table 18). Statistically significant (p=0.023 for the 100 mg and p<0.001 for the 300 mg dose) mean change from baseline in systolic blood pressure relative to placebo was -3.5 mmHg and -6 mmHg with canagliflozin 100 mg and 300 mg, respectively.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Fewer patients on canagliflozin in combination with metformin HCl and insulin required glycemic rescue therapy: 3.6% of patients receiving canagliflozin 100 mg, 2.7% of patients receiving canagliflozin 300 mg, and 6.2% of patients receiving placebo. An increased incidence of hypoglycemia was observed in this trial, which is consistent with the expected increase of hypoglycemia when an agent not associated with hypoglycemia is added to insulin [see *Warnings and Precautions (5.7) and Adverse Reactions (6.1)*].

Table 18: Results from 18–Week Placebo-Controlled Clinical Study of Canagliflozin in Combination with Metformin HCl and Insulin ≥ 30 Units/Day*

Efficacy Parameter	Placebo + Metformin HCl + Insulin (N=145)	Canagliflozin 100 mg + Metformin HCl + Insulin (N=139)	Canagliflozin 300 mg + Metformin HCl + Insulin (N=148)
HbA_{1c} (%)			
Baseline (mean)	8.15	8.20	8.22
Change from baseline (adjusted mean)	0.03	-0.64	-0.79
Difference from placebo (adjusted mean) (95% CI) [†]		-0.66 [‡] (-0.81, -0.51)	-0.82 [‡] (-0.96, -0.67)
Percent of patients achieving HbA _{1c} < 7%	9	19 [§]	29 [‡]
Fasting Plasma Glucose (mg/dL)			
Baseline	163	168	167
Change from baseline (adjusted mean)	1	-16	-24
Difference from placebo (adjusted mean) (97.5% CI) [†]		-16 [‡] (-28, -5)	-25 [‡] (-36, -14)
Body Weight			
Baseline (mean) in kg	102.3	99.7	101.1
% change from baseline (adjusted mean)	0.0	-1.7	-2.7
Difference from placebo (adjusted mean) (97.5% CI) [†]		-1.7 [‡] (-2.4, -1.0)	-2.7 [‡] (-3.4, -2.0)

* Intent-to-treat population using last observation in study prior to glycemic rescue therapy

[†] Least squares mean adjusted for baseline value and stratification factors

[‡] p<0.001

[§] p<0.01

14.2 Canagliflozin Cardiovascular Outcomes in Patients with Type 2 Diabetes Mellitus and Atherosclerotic Cardiovascular Disease

Canagliflozin is indicated to reduce the risk of major adverse cardiovascular events in adults with type 2 diabetes mellitus and established cardiovascular disease (CVD). However, the effectiveness of INVOKAMET/INVOKAMET XR on reducing the risk of major cardiovascular events in adults with type 2 diabetes and cardiovascular disease has not been established.

The CANVAS and CANVAS-R trials were multicenter, multi-national, randomized, double-blind parallel group, with similar inclusion and exclusion criteria. Patients eligible for enrollment in both CANVAS and CANVAS-R trials were: 30 years of age or older and had established, stable, cardiovascular, cerebrovascular, peripheral artery disease (66% of the enrolled population) or were 50 years of age or older and had two or more other specified risk factors for cardiovascular disease (34% of the enrolled population).

The integrated analysis of the CANVAS and CANVAS-R trials compared the risk of Major Adverse Cardiovascular Event (MACE) between canagliflozin and placebo when these were added to and used concomitantly with standard of care treatments for diabetes and atherosclerotic cardiovascular disease. The primary endpoint, MACE, was the time to first occurrence of a three-part composite outcome which included cardiovascular death, non-fatal myocardial infarction and non-fatal stroke.

In CANVAS, patients were randomly assigned 1:1:1 to canagliflozin 100 mg, canagliflozin 300 mg, or matching placebo. In CANVAS-R, patients were randomly assigned 1:1 to canagliflozin 100 mg or matching placebo, and titration to 300 mg was permitted at the investigator's discretion (based on tolerability and glycemic needs) after Week 13. Concomitant antidiabetic and atherosclerotic therapies could be adjusted, at the discretion of investigators, to ensure participants were treated according to the standard care for these diseases.

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

A total of 10,134 patients were treated (4,327 in CANVAS and 5,807 in CANVAS-R; total of 4,344 randomly assigned to placebo and 5,790 to canagliflozin) and exposed for a mean of 149 weeks (exposed for a mean of 223 weeks [4.3 years] in CANVAS and 94 weeks [1.8 years] in CANVAS-R). Approximately 78% of the trial population was Caucasian, 13% was Asian, and 3% was Black. The mean age was 63 years and approximately 64% were male.

The mean HbA_{1c} at baseline was 8.2% and mean duration of diabetes was 13.5 years with 70% of patients having had diabetes for 10 years or more. Approximately 31%, 21% and 17% reported a past history of neuropathy, retinopathy and nephropathy, respectively, and the mean eGFR 76 mL/min/1.73 m²). At baseline, patients were treated with one (19%) or more (80%) antidiabetic medications including metformin (77%), insulin (50%), and sulfonylurea (43%).

At baseline, the mean systolic blood pressure was 137 mmHg, the mean diastolic blood pressure was 78 mmHg, the mean LDL was 89 mg/dL, the mean HDL was 46 mg/dL, and the mean urinary albumin to creatinine ratio (UACR) was 115 mg/g. At baseline, approximately 80% of patients were treated with renin angiotensin system inhibitors, 53% with beta-blockers, 13% with loop diuretics, 36% with non-loop diuretics, 75% with statins, and 74% with antiplatelet agents (mostly aspirin). During the trial, investigators could modify anti-diabetic and cardiovascular therapies to achieve local standard of care treatment targets with respect to blood glucose, lipid, and blood pressure. More patients receiving canagliflozin compared to placebo initiated anti-thrombotics (5.2% vs 4.2%) and statins (5.8% vs 4.8%) during the trial.

For the primary analysis, a stratified Cox proportional hazards model was used to test for non-inferiority against a pre-specified risk margin of 1.3 for the hazard ratio of MACE.

In the integrated analysis of CANVAS and CANVAS-R trials, canagliflozin reduced the risk of first occurrence of MACE. The estimated hazard ratio (95% CI) for time to first MACE was 0.86 (0.75, 0.97). Refer to Table 19. Vital status was obtained for 99.6% of patients across the trials. The Kaplan-Meier curve depicting time to first occurrence of MACE is shown in Figure 3.

Table 19: Treatment Effect for the Primary Composite Endpoint, MACE, and its Components in the Integrated Analysis of CANVAS and CANVAS-R studies*

	Placebo N=4347	Canagliflozin N=5795	Hazard ratio (95% C.I.) [†]
Composite of cardiovascular death, non-fatal myocardial infarction, non-fatal stroke (time to first occurrence) ^{‡, §}	426 (10.4)	585 (9.2)	0.86 (0.75, 0.97)
Non-fatal myocardial infarction ^{‡, §}	159 (3.9)	215 (3.4)	0.85 (0.69, 1.05)
Non-fatal Stroke ^{‡, §}	116 (2.8)	158 (2.5)	0.90 (0.71, 1.15)
Cardiovascular Death ^{‡, §}	185 (4.6)	268 (4.1)	0.87 (0.72, 1.06)

* Intent-To-Treat Analysis Set


[†] P-value for superiority (2-sided) = 0.0158

[‡] Number and percentage of first events

[§] Due to pooling of unequal randomization ratios, Cochran-Mantel-Haenszel weights were applied to calculate percentages

[¶] Stratified Cox-proportional hazards model with treatment as a factor and stratified by study and by prior CV disease

Figure 3: Time to First Occurrence of MACE



INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

16 HOW SUPPLIED/STORAGE AND HANDLING

INVOKAMET tablets are available in bottles of 60 in the strengths listed below:

INVOKAMET	TABLET STRENGTH			
	50 mg/500 mg	50 mg/1000 mg	150 mg/500 mg	150 mg/1000 mg
Color	White	Beige	Yellow	Purple
Tablet Identification	CM	CM	CM	CM
	155	551	215	611
	Capsule-shaped, film-coated tablets			
NDC	50458-540-60	50458-541-60	50458-542-60	50458-543-60

INVOKAMET XR tablets are available in bottles of 60 in the strengths listed below:

INVOKAMET XR	STRENGTH			
	50 mg/500 mg	50 mg/1000 mg	150 mg/500 mg	150 mg/1000 mg
Color	Almost White to Light Orange	Pink	Orange	Reddish Brown
Tablet Identification	CM1	CM3	CM2	CM4
	Oblong, biconvex, film-coated tablets, a thin line on the tablet side may be visible.			
NDC	50458-940-01	50458-941-01	50458-942-01	50458-943-01

Storage and Handling

Keep out of reach of children.

Store at 68-77°F (20-25°C); excursions permitted between 59°F and 86°F (15°C and 30°C) [see USP Controlled Room Temperature]. Store and dispense in the original container. Storage in a pill box or pill organizer is allowed for up to 30 days.

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-Approved Patient Labeling (Medication Guide).

Lactic Acidosis:

Explain the risks of lactic acidosis, its symptoms, and conditions that predispose to its development, as noted in *Warnings and Precautions (5.1)*. Advise patients to discontinue INVOKAMET/INVOKAMET XR immediately and to promptly notify their healthcare provider if unexplained hyperventilation, myalgias, malaise, unusual somnolence or other nonspecific symptoms occur. Once a patient is stabilized on INVOKAMET/INVOKAMET XR, gastrointestinal symptoms, which are common during initiation of metformin, are unlikely to recur. Later occurrence of gastrointestinal symptoms could be due to lactic acidosis or other serious disease.

Counsel patients against excessive alcohol intake while receiving INVOKAMET/INVOKAMET XR.

Inform patients about importance of regular testing of renal function and hematological parameters while receiving INVOKAMET/INVOKAMET XR.

Instruct patients to inform their doctor that they are taking INVOKAMET/INVOKAMET XR prior to any surgical or radiological procedure, as temporary discontinuation of INVOKAMET/INVOKAMET XR may be required until renal function has been confirmed to be normal [see *Warnings and Precautions (5.1)*].

Lower Limb Amputation:

Inform patients that INVOKAMET/INVOKAMET XR is associated with an increased risk of amputations. Counsel patients about the importance of routine preventative foot care. Instruct patients to monitor for new pain or tenderness, sores or ulcers, or infections involving the leg or foot and to seek medical advice immediately if such signs or symptoms develop [see *Boxed Warning and Warnings and Precautions (5.2)*].

Hypotension:

Inform patients that symptomatic hypotension may occur with INVOKAMET/INVOKAMET XR and advise them to contact their doctor if they experience such symptoms [see *Warnings and Precautions (5.3)*]. Inform patients that dehydration may increase the risk for hypotension and to have adequate fluid intake.

Ketoacidosis:

Inform patients that ketoacidosis is a serious life-threatening condition. Cases of ketoacidosis have been reported during use of canagliflozin. Instruct patients to check ketones (when possible) if symptoms consistent with ketoacidosis occur even if blood glucose is not elevated. If symptoms of ketoacidosis (including nausea, vomiting, abdominal pain, tiredness, and labored breathing) occur, instruct patients to discontinue INVOKAMET/INVOKAMET XR and seek medical advice immediately [see *Warnings and Precautions (5.4)*].

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Acute Kidney Injury:

Inform patients that acute kidney injury has been reported during use of canagliflozin. Advise patients to seek medical advice immediately if they have reduced oral intake (such as due to acute illness or fasting), or increased fluid losses (such as due to vomiting, diarrhea, or excessive heat exposure), as it may be appropriate to temporarily discontinue INVOKAMET/INVOKAMET XR use in those settings [see *Warnings and Precautions (5.5)*].

Serious Urinary Tract Infections:

Inform patients of the potential for urinary tract infections, which may be serious. Provide them with information on the symptoms of urinary tract infections. Advise them to seek medical advice if such symptoms occur [see *Warnings and Precautions (5.6)*].

Necrotizing Fasciitis of the Perineum (Fournier's Gangrene):

Inform patients that necrotizing infections of the perineum (Fournier's gangrene) have occurred with INVOKAMET/INVOKAMET XR. Counsel patients to promptly seek medical attention if they develop pain or tenderness, redness, or swelling of the genitals or the area from the genitals back to the rectum, along with a fever above 100.4°F or malaise [see *Warnings and Precautions (5.8)*].

Genital Mycotic Infections in Females (e.g., Vulvovaginitis):

Inform female patients that vaginal yeast infection (e.g., vulvovaginitis) may occur and provide them with information on the signs and symptoms of a vaginal yeast infection. Advise them of treatment options and when to seek medical advice [see *Warnings and Precautions (5.9)*].

Genital Mycotic Infections in Males (e.g., Balanitis or Balanoposthitis):

Inform male patients that yeast infection of penis (e.g., balanitis or balanoposthitis) may occur, especially in uncircumcised males and patients with prior history. Provide them with information on the signs and symptoms of balanitis and balanoposthitis (rash or redness of the glans or foreskin of the penis). Advise them of treatment options and when to seek medical advice [see *Warnings and Precautions (5.9)*].

Hypersensitivity Reactions:

Inform patients that serious hypersensitivity reactions, such as urticaria, rash, anaphylaxis, and angioedema, have been reported with canagliflozin. Advise patients to report immediately any signs or symptoms suggesting allergic reaction and to discontinue drug until they have consulted prescribing physicians [see *Warnings and Precautions (5.10)*].

Bone Fracture:

Inform patients that bone fractures have been reported in patients taking canagliflozin. Provide them with information on factors that may contribute to fracture risk [see *Warnings and Precautions (5.11)*].

Laboratory Tests:

Inform patients that they will test positive for glucose in their urine while on INVOKAMET/INVOKAMET XR [see *Drug Interactions (7.2)*].

Females of Reproductive Age:

Advise pregnant women, and females of reproductive potential of the potential risk to a fetus with treatment with INVOKAMET/INVOKAMET XR [see *Use in Specific Populations (8.1)*]. Instruct females of reproductive potential to report pregnancies to their physicians as soon as possible.

Inform females that treatment with INVOKAMET/INVOKAMET XR may result in ovulation in some premenopausal anovulatory women which may lead to unintended pregnancy [see *Use in Specific Populations (8.3)*].

Lactation:

Advise women that breastfeeding is not recommended during treatment with INVOKAMET/INVOKAMET XR [see *Use in Specific Populations (8.2)*].

INVOKAMET® (canagliflozin and metformin hydrochloride) tablets
INVOKAMET® XR (canagliflozin and metformin hydrochloride) extended-release tablets

Administration:

Instruct patients to keep INVOKAMET/INVOKAMET XR in the original bottle to protect from moisture. Advise patients that storage in a pill box or pill organizer is allowed for up to 30 days.

Instruct patients to take INVOKAMET only as prescribed twice daily with food. If a dose is missed, advise patients not to take two doses of INVOKAMET at the same time.

Instruct patients to take INVOKAMET XR only as prescribed once daily with the morning meal. If a dose is missed, advise patients to take it as soon as it is remembered unless it is almost time for the next dose, in which case patients should skip the missed dose and take the medicine at the next regularly scheduled time. Advise patients not to take more than two tablets of INVOKAMET XR at the same time.

Instruct patients that INVOKAMET XR must be swallowed whole and never crushed, cut, or chewed, and that the inactive ingredients may occasionally be eliminated in the feces as a soft mass that may resemble the original tablet.

Manufactured for:
Janssen Pharmaceuticals, Inc.
Titusville, NJ 08560

INVOKAMET

Finished product manufactured by:
Janssen Ortho LLC
Gurabo, PR 00778

Or

Janssen Cilag SpA
Latina, Italy

INVOKAMET XR

Finished product manufactured by:
Janssen Ortho LLC
Gurabo, PR 00778

Licensed from Mitsubishi Tanabe Pharma Corporation

© 2014, 2016 Janssen Pharmaceutical Companies

Medication Guide
INVOKAMET® (in vok' a met)
(canagliflozin and metformin hydrochloride) tablets, for oral use
and
INVOKAMET® (in vok' a met) XR
(canagliflozin and metformin hydrochloride) extended-release tablets, for oral use

What is the most important information I should know about INVOKAMET and INVOKAMET XR?

INVOKAMET and INVOKAMET XR can cause serious side effects, including:

- **Lactic Acidosis.** Metformin, one of the medicines in INVOKAMET and INVOKAMET XR, can cause a rare but serious condition called lactic acidosis (a build-up of lactic acid in the blood) that can cause death. Lactic acidosis is a medical emergency and must be treated in the hospital.

Call your doctor right away if you have any of the following symptoms, which could be signs of lactic acidosis:

- you feel cold in your hands or feet
- you feel very weak or tired
- you have trouble breathing
- you have stomach pains, nausea, or vomiting
- you have a slow or irregular heartbeat
- you have unusual (not normal) muscle pain
- you have unusual sleepiness or sleep longer than usual
- you feel dizzy or lightheaded

Most people who have had lactic acidosis had other conditions that, in combination with metformin use, led to the lactic acidosis. Tell your doctor if you have any of the following, because you have a higher chance for getting lactic acidosis with INVOKAMET or INVOKAMET XR if you:

- have severe kidney problems or your kidneys are affected by certain x-ray tests that use injectable dye.
- have liver problems.
- drink alcohol very often, or drink a lot of alcohol in short-term “binge” drinking.
- get dehydrated (lose a large amount of body fluids). This can happen if you are sick with a fever, vomiting, or diarrhea. Dehydration can also happen when you sweat a lot with activity or exercise and do not drink enough fluids.
- have surgery.
- have a heart attack, severe infection, or stroke.

The best way to keep from having a problem with lactic acidosis from metformin is to tell your doctor if you have any of the problems in the list above. Your doctor will decide to stop your INVOKAMET or INVOKAMET XR for a while if you have any of these things.

- **Amputations.** INVOKAMET or INVOKAMET XR may increase your risk of lower limb amputations. Amputations mainly involve removal of the toe or part of the foot, however, amputations involving the leg, below and above the knee, have also occurred. Some people had more than one amputation, some on both sides of the body.

You may be at a higher risk of lower limb amputation if you:

- have a history of amputation
- have heart disease or are at risk for heart disease
- have had blocked or narrowed blood vessels, usually in your leg
- have damage to the nerves (neuropathy) in your leg
- have had diabetic foot ulcers or sores

Call your doctor right away if you have new pain or tenderness, any sores, ulcers, or infections in your leg or foot. Your doctor may decide to stop your INVOKAMET or INVOKAMET XR for a while if you have any of these signs or symptoms.

Talk to your doctor about proper foot care.

INVOKAMET or INVOKAMET XR can have other serious side effects. See “What are the possible side effects of INVOKAMET or INVOKAMET XR?”

What is INVOKAMET or INVOKAMET XR?

- INVOKAMET contains 2 prescription medicines called canagliflozin (INVOKANA) and metformin hydrochloride (GLUCOPHAGE). INVOKAMET XR contains 2 prescription medicines called canagliflozin (INVOKANA) and metformin hydrochloride extended-release (GLUMETZA). INVOKAMET or INVOKAMET XR can be used:
 - along with diet and exercise to lower blood sugar (glucose) in adults with type 2 diabetes
- in adults with type 2 diabetes who have known cardiovascular disease when both canagliflozin and metformin are appropriate and canagliflozin is needed to reduce the risk of major cardiovascular events such as heart attack, stroke, or death.
- INVOKAMET or INVOKAMET XR is not for people with type 1 diabetes.
- INVOKAMET or INVOKAMET XR is not for people with diabetic ketoacidosis (increased ketones in blood or urine).
- It is not known if INVOKAMET or INVOKAMET XR is safe and effective in children under 18 years of age.

Who should not take INVOKAMET or INVOKAMET XR?

Do not take INVOKAMET or INVOKAMET XR if you:

- have moderate to severe kidney problems or are on dialysis.
- have a condition called metabolic acidosis or diabetic ketoacidosis (increased ketones in the blood or urine).
- are allergic to canagliflozin, metformin, or any of the ingredients in INVOKAMET or INVOKAMET XR. See the end of this Medication Guide for a list of ingredients in INVOKAMET and INVOKAMET XR. Symptoms of allergic reaction to INVOKAMET and INVOKAMET XR may include:
 - rash
 - raised red patches on your skin (hives)
 - swelling of the face, lips, mouth, tongue, and throat that may cause difficulty in breathing or swallowing

What should I tell my doctor before taking INVOKAMET or INVOKAMET XR?

Before you take INVOKAMET or INVOKAMET XR, tell your doctor if you:

- have a history of amputation.
- have heart disease or are at risk for heart disease.
- have had blocked or narrowed blood vessels, usually in your leg.
- have damage to the nerves (neuropathy) in your leg.
- have had diabetic foot ulcers or sores.
- have moderate to severe kidney problems.
- have liver problems.
- have a history of urinary tract infections or problems with urination.
- are on a low sodium (salt) diet. Your doctor may change your diet or your dose of INVOKAMET or INVOKAMET XR.
- have ever had an allergic reaction to INVOKAMET or INVOKAMET XR.
- are going to get an injection of dye or contrast agents for an x-ray procedure. INVOKAMET or INVOKAMET XR may need to be stopped for a short time. Talk to your doctor about when you should stop INVOKAMET or INVOKAMET XR and when you should start INVOKAMET or INVOKAMET XR again. See **“What is the most important information I should know about INVOKAMET or INVOKAMET XR?”**
- have heart problems, including congestive heart failure.
- are going to have surgery.
- are eating less due to illness, surgery, or a change in your diet.
- have or have had problems with your pancreas, including pancreatitis or surgery on your pancreas.
- drink alcohol very often, or drink a lot of alcohol in the short-term (“binge” drinking).
- have other medical conditions.
- are pregnant or plan to become pregnant. INVOKAMET or INVOKAMET XR may harm your unborn baby. If you become pregnant while taking INVOKAMET or INVOKAMET XR, tell your doctor as soon as possible. Talk with your doctor about the best way to control your blood sugar while you are pregnant.
- are a premenopausal woman (before the “change of life”), who does not have periods regularly or at all. INVOKAMET or INVOKAMET XR may increase your chance of becoming pregnant. Talk to your doctor about birth control choices while taking INVOKAMET or INVOKAMET XR, if you are not planning to become pregnant. Tell your doctor right away if you become pregnant while taking INVOKAMET or INVOKAMET XR.
- are breastfeeding or plan to breastfeed. INVOKAMET or INVOKAMET XR may pass into your breast milk and may harm your baby. Talk with your doctor about the best way to feed your baby if you are taking INVOKAMET or INVOKAMET XR. Do not breastfeed while taking INVOKAMET or INVOKAMET XR.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

INVOKAMET or INVOKAMET XR may affect the way other medicines work and other medicines may affect how INVOKAMET or INVOKAMET XR works. Especially tell your doctor if you take:

- diuretics (water pills)
- phenytoin or phenobarbital (used to control seizures)
- digoxin (Lanoxin®)* (used to treat heart problems)
- rifampin (used to treat or prevent tuberculosis)
- ritonavir (Norvir®, Kaletra®)* (used to treat HIV infection)

Ask your doctor or pharmacist for a list of these medicines if you are not sure if your medicine is listed above.

Know the medicines you take. Keep a list of them and show it to your doctor and pharmacist when you get a new medicine.

How should I take INVOKAMET or INVOKAMET XR?

- If you are prescribed INVOKAMET, take by mouth 2 times each day with meals exactly as your doctor tells you to take it. Taking INVOKAMET with meals may lower your chance of having an upset stomach.
- If you are prescribed INVOKAMET XR, take by mouth 1 time each day with the morning meal exactly as your doctor tells you to take it. Taking INVOKAMET XR with a meal may lower your chance of having an upset stomach.
- Swallow INVOKAMET XR whole. Do not crush, cut, or chew.
- You may sometimes pass a soft mass in your stools (bowel movement) that looks like INVOKAMET XR tablets. It is normal to see this in your stool.
- Your doctor will tell you how much INVOKAMET or INVOKAMET XR to take and when to take it. Your doctor may change your dose if needed.
- Your doctor may tell you to take INVOKAMET or INVOKAMET XR along with other diabetes medicines. Low blood sugar can happen more often when INVOKAMET or INVOKAMET XR is taken with certain other diabetes medicines. See **“What are the possible side effects of INVOKAMET or INVOKAMET XR?”**
- If you miss a dose of INVOKAMET, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at the next regularly scheduled time. Do not take 2 tablets of INVOKAMET at the same time. Talk to your doctor if you have questions about a missed dose.
- If you miss a dose of INVOKAMET XR, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose and take the medicine at the next regularly scheduled time. Do not take more than 2 tablets of INVOKAMET XR at the same time. Talk to your doctor if you have questions about a missed dose.
- If you take too much INVOKAMET or INVOKAMET XR, call your doctor or go to the nearest hospital emergency room right away.
- When your body is under some types of stress, such as fever, trauma (such as a car accident), infection, or surgery, the amount of diabetes medicine you need may change. Tell your doctor right away if you have any of these conditions and follow your doctor’s instructions.
- Stay on your prescribed diet and exercise program while taking INVOKAMET or INVOKAMET XR.
- Check your blood sugar as your doctor tells you to.
- INVOKAMET and INVOKAMET XR will cause your urine to test positive for glucose.
- Your doctor may do certain blood tests before you start INVOKAMET or INVOKAMET XR and during treatment as needed. Your doctor may change your dose of INVOKAMET or INVOKAMET XR based on the results of your blood tests.
- Your doctor will check your diabetes with regular blood tests, including your blood sugar levels and your hemoglobin A_{1c}.

What should I avoid while taking INVOKAMET or INVOKAMET XR?

- Avoid drinking alcohol very often, or drinking a lot of alcohol in a short period of time (“binge” drinking). It can increase your chances of getting serious side effects.

What are the possible side effects of INVOKAMET or INVOKAMET XR?

INVOKAMET or INVOKAMET XR may cause serious side effects including:

- See **“What is the most important information I should know about INVOKAMET or INVOKAMET XR?”**
- **dehydration.** INVOKAMET or INVOKAMET XR can cause some people to become dehydrated (the loss of too much body water). Dehydration may cause you to feel dizzy, faint, lightheaded, or weak, especially when you stand up (orthostatic hypotension).

You may be at higher risk of dehydration if you:

- have low blood pressure
- take medicines to lower your blood pressure, including diuretics (water pill)
- are on a low sodium (salt) diet
- have kidney problems
- are 65 years of age or older

Talk to your doctor about what you can do to prevent dehydration including how much fluid you should drink on a daily basis.

- **ketoacidosis (increased ketones in your blood or urine).** Ketoacidosis has happened in people who have **type 1 diabetes or type 2 diabetes**, during treatment with canagliflozin, one of the medicines in INVOKAMET and INVOKAMET XR. Ketoacidosis is a serious condition, which may need to be treated in a hospital. Ketoacidosis may lead to death. **Ketoacidosis can happen with INVOKAMET or INVOKAMET XR, even if your blood sugar is less than 250 mg/dL. Stop taking INVOKAMET or INVOKAMET XR and call your doctor right away if you get any of the following symptoms:**

- nausea
- vomiting
- stomach area (abdominal) pain
- tiredness
- trouble breathing

What are the possible side effects of INVOKAMET or INVOKAMET XR? (continued)

If you get any of these symptoms during treatment with INVOKAMET or INVOKAMET XR, if possible, check for ketones in your urine, even if your blood sugar is less than 250 mg/dL.

- **kidney problems.** Sudden kidney injury has happened to people taking INVOKAMET or INVOKAMET XR. Talk to your doctor right away if you:
 - reduce the amount of food or liquid you drink for example, if you are sick or cannot eat or
 - you start to lose liquids from your body for example, from vomiting, diarrhea or being in the sun too long.
- **serious urinary tract infections.** Serious urinary tract infections that may lead to hospitalization have happened in people who are taking canagliflozin, one of the medicines in INVOKAMET and INVOKAMET XR. Tell your doctor if you have any signs or symptoms of a urinary tract infection such as a burning feeling when passing urine, a need to urinate often, the need to urinate right away, pain in the lower part of your stomach (pelvis), or blood in the urine. Sometimes people may also have a fever, back pain, nausea, or vomiting.
- **low blood sugar (hypoglycemia).** If you take INVOKAMET or INVOKAMET XR with another medicine that can cause low blood sugar, such as a sulfonyleurea or insulin, your risk of getting low blood sugar is higher. The dose of your sulfonyleurea medicine or insulin may need to be lowered while you take INVOKAMET or INVOKAMET XR. Signs and symptoms of low blood sugar may include:
 - headache
 - drowsiness
 - weakness
 - confusion
 - dizziness
 - irritability
 - hunger
 - fast heartbeat
 - sweating
 - shaking or feeling jittery
- **a rare but serious bacterial infection that causes damage to the tissue under the skin (necrotizing fasciitis) in the area between and around the anus and genitals (perineum).** Necrotizing fasciitis of the perineum has happened in women and men who take canagliflozin, one of the medicines in INVOKAMET and INVOKAMET XR. Necrotizing fasciitis of the perineum may lead to hospitalization, may require multiple surgeries, and may lead to death. **Seek medical attention immediately if you have fever or you are feeling very weak, tired or uncomfortable (malaise) and you develop any of the following symptoms in the area between and around your anus and genitals:**
 - pain or tenderness
 - swelling
 - redness of the skin (erythema)
- **vaginal yeast infection.** Women who take INVOKAMET or INVOKAMET XR may get vaginal yeast infections. Symptoms of a vaginal yeast infection include:
 - vaginal odor
 - white or yellowish vaginal discharge (discharge may be lumpy or look like cottage cheese)
 - vaginal itching
- **yeast infection of the penis (balanitis or balanoposthitis).** Men who take INVOKAMET or INVOKAMET XR may get a yeast infection of the skin around the penis. Certain men who are not circumcised may have swelling of the penis that makes it difficult to pull back the skin around the tip of the penis. Other symptoms of yeast infection of the penis include:
 - redness, itching, or swelling of the penis
 - rash of the penis
 - foul smelling discharge from the penis
 - pain in the skin around the penis

Talk to your doctor about what to do if you get symptoms of a yeast infection of the vagina or penis. Your doctor may suggest you use an over-the-counter antifungal medicine. Talk to your doctor right away if you use an over-the-counter antifungal medication and your symptoms do not go away.

- **serious allergic reaction.** If you have any symptoms of a serious allergic reaction, stop taking INVOKAMET or INVOKAMET XR and call your doctor right away or go to the nearest hospital emergency room. See **“Who should not take INVOKAMET or INVOKAMET XR?”**. Your doctor may give you a medicine for your allergic reaction and prescribe a different medicine for your diabetes.
- **broken bones (fractures).** Bone fractures have been seen in patients taking canagliflozin. Talk to your doctor about factors that may increase your risk of bone fracture.
- **low vitamin B₁₂ (vitamin B₁₂ deficiency).** Using metformin for long periods of time may cause a decrease in the amount of vitamin B₁₂ in your blood, especially if you have had low vitamin B₁₂ blood levels before. Your doctor may do blood tests to check your vitamin B₁₂ levels.

Other common side effects of INVOKAMET or INVOKAMET XR include:

- nausea and vomiting
- diarrhea
- weakness
- gas
- upset stomach
- indigestion
- headache
- changes in urination, including urgent need to urinate more often, in larger amounts, or at night

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of INVOKAMET or INVOKAMET XR. For more information, ask your doctor or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

You may also report side effects to Janssen Pharmaceuticals, Inc. at 1-800-526-7736.

How should I store INVOKAMET or INVOKAMET XR?

- Store INVOKAMET and INVOKAMET XR at room temperature between 68°F to 77°F (20°C to 25°C).
- Store in the original container to protect from moisture. Storage in a pill box or pill organizer is allowed for up to 30 days.

Keep INVOKAMET and INVOKAMET XR and all medicines out of the reach of children.

General information about the safe and effective use of INVOKAMET and INVOKAMET XR.

Medicines are sometimes prescribed for purposes other than those listed in the Medication Guide. Do not use INVOKAMET or INVOKAMET XR for a condition for which it was not prescribed. Do not give INVOKAMET or INVOKAMET XR to other people, even if they have the same symptoms you have. It may harm them.

This Medication Guide summarizes the most important information about INVOKAMET and INVOKAMET XR. If you would like more information, talk with your doctor. You can ask your pharmacist or doctor for information about INVOKAMET or INVOKAMET XR that is written for healthcare professionals.

For more information about INVOKAMET or INVOKAMET XR, call 1-800-526-7736 or visit our websites at www.invokamet.com or www.invokametxr.com.

What are the ingredients of INVOKAMET?

Active ingredients: canagliflozin and metformin hydrochloride

Inactive ingredients: The tablet core contains croscarmellose sodium, hypromellose, magnesium stearate, and microcrystalline cellulose. The magnesium stearate is vegetable-sourced. In addition, the tablet coating contains Macrogol/PEG3350, polyvinyl alcohol (partially hydrolyzed), talc, titanium dioxide, iron oxide yellow (50 mg/1,000 mg and 150 mg/500 mg tablets only), iron oxide red (50 mg/1,000 mg, 150 mg/500 mg and 150 mg/1,000 mg tablets only), and iron oxide black (150 mg/1,000 mg tablets only).

What are the ingredients of INVOKAMET XR?

Active ingredients: canagliflozin and metformin hydrochloride

Inactive ingredients: The tablet core contains croscarmellose sodium, hydroxypropyl cellulose, hypromellose, lactose anhydrous, magnesium stearate (vegetable-sourced), microcrystalline cellulose, polyethylene oxide, and silicified microcrystalline cellulose (50 mg/500 mg and 50 mg/1,000 mg tablets only). In addition, the tablet coating contains macrogol/PEG3350, polyvinyl alcohol (partially hydrolyzed), talc, titanium dioxide, iron oxide red, iron oxide yellow, and iron oxide black (50 mg/1,000 mg and 150 mg/1,000 mg tablets only).

*The brands listed are trademarks of their respective owners and are not trademarks of Janssen Pharmaceuticals, Inc.

Manufactured for: Janssen Pharmaceuticals, Inc., Titusville, NJ 08560. INVOKAMET manufactured by: Janssen Ortho LLC, Gurabo, PR 00778 or Janssen Cilag SpA, Latina, Italy. INVOKAMET XR manufactured by: Janssen Ortho LLC, Gurabo, PR 00778. Licensed from Mitsubishi Tanabe Pharma Corporation. © 2014, 2016 Janssen Pharmaceutical Companies

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Revised 10/2018

cp-54215v3